



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/45, 15/62, C07K 14/115, 14/12, 14/135, A61K 39/155, 39/165</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/18929</b> (43) International Publication Date: <b>6 April 2000 (06.04.00)</b>
(21) International Application Number: <b>PCT/EP99/07004</b> (22) International Filing Date: <b>20 September 1999 (20.09.99)</b> (30) Priority Data: 9820931.5      25 September 1998 (25.09.98)      GB 9906868.6      24 March 1999 (24.03.99)      GB (71) Applicant (for all designated States except US): <b>SMITHKLINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).</b> (72) Inventors; and (75) Inventors/Applicants (for US only): <b>BOLLEN, Alex [BE/BE]; Université Libre de Bruxelles, Faculté des Sciences, Rue de l'Industrie 24, B-1440 Nivelles (BE). HOUARD, Sophie [BE/BE]; CRI - Centre de Recherches Industriel/ULB, Département de Biologie Moléculaire, Rue de l'Industrie 24, B-1400 Nivelles (BE).</b> (74) Agent: <b>PRIVETI, Kathryn, Louise; Corporate Intellectual Property, SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).</b>			(81) Designated States: <b>AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: <b>NOVEL COMPOUNDS</b> (57) Abstract  Heterochimeric proteins or immunogenic derivatives thereof are described comprising immunogenic fragments of RSV, PIV1, PIV2, PIV3, MV and MuV fusion and attachment glycoproteins. Such heterochimeric proteins may be expressed, in particular, in CHO cells and may be used in vaccine compositions to treat respiratory disorders such as those caused by paramyxoviridae viral antigens.			

ATTORNEY DOCKET NUMBER: 7682-059-999  
SERIAL NUMBER: 09/923,070  
REFERENCE: **CI**

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## Novel Compounds

The present invention relates to recombinant heterochimeric paramyxoviridae glycoproteins and their expression in eukaryotic cells, particularly in Chinese Hamster Ovary (CHO) cells. The invention further relates to methods for constructing and expressing such heterochimeric proteins, intermediates for use therein, methods to optimize the codon usage of the nucleic acid sequences which encode such heterochimeric proteins and the use of the recombinant proteins as vaccines for the prevention of diseases caused by paramyxoviridae pathogens.

10

The mumps (MuV), Measles (MV), the parainfluenza type I (PIV1), type II (PIV2) and type III (PIV3) and the respiratory syncytial (RSV) virus belong to the paramyxoviridae family. The MuV is classified in the rubulavirus subclass, the MV is classified in the Morbillivirus subclass, the parainfluenza viruses (PIV1, PIV2 and PIV3) are classified in the paramyxovirus subclass while the RSV is attached to the pneumovirus subclass.

15

RSV is the most important cause of viral lower respiratory tract disease in infants and children. The fusion (F) and the attachment (G) protein which are both viral surface glycoproteins appear to be of potential value for the development of a vaccine against RSV.

20

The fusion protein F of RSV contains 574 amino acid residues; amino acids 1 to 21 correspond to the signal peptide and residues 525 to 549 to the membrane anchor domain. The molecule presents five potential sites for glycosylation. The F protein is synthesized as a 70 kDa precursor ( $F_0$ ) which undergoes proteolytic maturation to yield the  $F_1$  subunit (48 kDa) and  $F_2$  (23 kDa) linked via disulfide bridges. The protein F, when injected into animals, leads to the production of neutralizing antibodies and may induce cytotoxic lymphocytes (CTLs).

25

30

The attachment or G protein of RSV contains 298 amino acid residues and is heavily glycosylated since half of its molecular mass (90 kDa) is contributed by

oligosaccharide side chains, chiefly in the form of O-linked sugars. It has been shown that the G protein, when injected into animals, provides protection against homologous but not heterologous subgroup virus challenge. This protein is extremely variable and there is only a stretch of 13 amino acid residues which is conserved in all RSV.

The PIV3 is second to RSV as a major agent of severe viral respiratory tract infections in infants. The fusion protein F of PIV3 contains 539 amino acid residues; amino acids 1 to 18 correspond to the signal peptide and residues 494 to 516 to the membrane anchor domain. The molecule presents 4 potential sites for glycosylation. The F protein is synthesized as a 70 kDa precursor ( $F_0$ ) which undergoes proteolytic maturation to yield the  $F_1$  (56 kDa) and  $F_2$  (14 kDa) subunits linked via disulfide bridges. The protein  $F_1$ , when injected into animals, leads to the production of neutralizing antibodies. The F protein is involved in cell fusion during viral infection and carries an hemolysin activity. Used alone for immunization, the F protein generates an immune response which is insufficient to confer protection against a challenge with the virus. Complete protection is only acquired by concomitant immunization with the attachment protein HN, another glycoprotein of PIV3.

The protein HN carries hemagglutinin and neuraminidase activities. It is composed of 572 amino acids; its membrane anchor domain occurs in the N-terminal end of the molecule, between amino acid residues 32 and 53. Four potential sites for glycosylation have been identified. Injection of protein HN into animals generates an immune response and neutralizing antibodies. These antibodies however do not protect completely against a challenge with the virus. Full protection is obtained only by concomitant immunization with the F protein of PIV3.

The PIV1 virus was initially isolated from young children suffering from disorders of the lower respiratory tract. Infection with PIV1 causes the majority of cases of croup found for all infections caused by paramyxoviruses. Viral transmission of

PIV1 is by person to person contact or by aerosol, although the virus does not persist in the environment for long.

5 Like PIV2 and PIV3, the PIV1 virus has two surface glycoproteins, the fusion protein (F) and the attachment protein (HN). These two proteins are the priority targets for the development of a subunit vaccine, the properties of which would be to ensure protection of children from the very first months of life and to prevent reinfection, or at least to prevent the serious complications by restricting viral development to the upper respiratory tract where the consequences would be benign  
10 (common cold).

PIV2 also affects very young children and causes the same type of respiratory disorders, essentially croup, but of less severity. The PIV2 virus has two surface glycoproteins (F and HN), which are potential targets for the development of a  
15 subunit vaccine.

The measles virus is an extremely contagious agent which establishes itself in the epithelial cells of the respiratory tract, the oropharynx or the conjunctiva. The infection causes fever, cough, head-cold, conjunctivitis and a characteristic  
20 generalised rash.

There is no appropriate inactivated vaccine against measles but an effective attenuated live vaccine is available and is generally used in combination with the attenuated live vaccines against rubella and mumps. This live vaccine protects  
25 against the disease for at least 20 years. The measles virus has two surface glycoproteins, which are potential targets for the development of a subunit vaccine. The fusion protein (F) is a 550 amino acid long glycosylated molecule and, as for the other paramyxovirus, has to undergo proteolytic cleavage to yield F<sub>1</sub> and F<sub>2</sub> subunits that are linked via disulfide bridges. This molecule, which carries a  
30 haemolysin activity, generates an immune protective response when injected into animals. The attachment protein (H), is a 617 amino acid long glycosylated protein, which carries a hemagglutinin activity. This protein leads, when injected into

animals, to the production of neutralizing antibodies that are able to inhibit hemagglutination. This immune response protects the animal against a viral challenge.

- 5 The mumps virus is a pathogen causing the contagious infantile illness which consists of the inflammation of parotid glands. During the incubation period following infection, the virus replicates in the respiratory epithelium then disseminates into secretory ducts of the parotid glands. Other glands may become infected thereafter and numerous cases of meningitis have been reported. Among  
10 complications related to the infection, encephalitis is a serious one, with a mortality rate of about 1%; deafness cases have also been reported.

- A vaccine against mumps is available: it is made of an attenuated live virus, produced by culturing infected embryonic chicken cells. The vaccine leads to the  
15 seroconversion in vaccinated individuals and protects against infection in more than 95% of seronegative persons. The vaccine thus reduces significantly the frequencies of complications.

- In a number of cases, however, viral infection is not detected because the effects  
20 remain subclinical. Young children and aged people are most likely to develop complications from mumps infection. In view of the inherent risks related to the use of attenuated live vaccines, such as the potentiation of the illness upon natural surinfection in vaccinated individuals, it is desirable to improve the safety of the vaccine, particularly for the groups at risk.

- 25 The fusion protein F of mumps virus contains 538 amino acid residues; amino acids 1 to 26 correspond to the signal peptide and residues 483 to 512 to the membrane anchor domain. The molecule presents 7 potential sites for glycosylation. The F protein is synthesized as a 65-74 kDa precursor ( $F_0$ ) which undergoes proteolytic  
30 maturation to yield the  $F_1$  (58-61 kDa) and  $F_2$  (10-16 kDa) subunits linked via disulfide bridges. The protein F is involved in cell fusion during viral infection, carries an haemolysin activity and plays a role for viral penetration into cells. It

does not however carry the antibody dependent cellular cytotoxicity (ADCC) as observed for another mumps virus glycoprotein, HN.

5 The protein HN (molecular weight 74-80 kDa) carries hemagglutinin and neuraminidase activities which are involved in virus attachment to cells and in the disruption of the host cell membranes. Protein HN (attachment protein or hemagglutinin-neuraminidase) generates neutralizing antibodies and appears important for the development of ADCC. Protein HN is composed of 582 amino acids; it carries a N-terminal anchor domain (residues 33 to 52) and 9 potential sites  
10 for glycosylation.

For the viruses considered above, it appears that concomitant immunization with both membrane glycoproteins F and HN, or G in the case of RSV, are required to achieve full protection in the animal model. Chimeric proteins containing both the F  
15 and G proteins of RSV, or the F and HN proteins of PIV3 have shown complete protection against RSV or PIV3 challenge in cotton rats (Brideau et al, J Gen Virol, 1989, 70 2637-2644 and Brideau et al, J Gen Virol, 1993, 74, 471-477).

WO9314207 (Connaught) describes heterochimeric proteins comprising RSV and  
20 PIV3 proteins including F(RSV)xHN(PIV3) and F(PIV3)xG(RSV) hybrids, and suggests that such proteins can be expressed from a variety of host cells including bacterial, mammalian, insect, yeast and fungal cells. The specific examples describe expression in insect Sf9 and High 5 cells and mammalian Vero cells. There is no specific disclosure of the use of CHO cells. The use of Sf9 and High 5  
25 cells is also described by Du et al, BIO/TECHNOLOGY 12, 1994, 813-818.

Homa et al (Upjohn), J Gen Virol, 1993, 74, 1995-1999 describes another heterochimeric protein, F(RSV)xHN(PIV3) expressed in insect cells using a recombinant baculovirus.

30

Homochimeric paramyxoviridae glycoproteins have also been described by several workers:-

WO8905823 (Upjohn) describes RSV FxG and GxF hybrids which can be expressed from bacterial, yeast, mammalian and insect cells. Example 7 describes the expression of an RSV FxG protein from CHO cells although there are no details of  
5 how successful such expression is.

WO8910405 (Upjohn) describes PIV3 FxHN and HNxF hybrids which can be expressed from bacterial, yeast, mammalian and insect cells. Example 6 describes the expression of a PIV3 FxHN protein from CHO cells, however no details are  
10 given quantifying the extent of expression and secretion.

Lehman et al (Upjohn), J Gen Virol, 1993, 74, 459-469 describes the expression of PIV3 FxHN in insect cells using recombinant baculovirus vectors as well as in CHO cells.

15 WO9306218 (SmithKline Beecham Biologicals) describes PIV3 FxHN hybrids which can be expressed in eukaryotic cells including vaccinia, CHO or Vero cells. Example B)2 describes the expression of a  $Fs^+a^+xHN^+$  hybrid in CHO cells and indicates that the product was almost evenly distributed between cells and medium.  
20 No details are however given quantifying the extent of expression and secretion.

WO9425600 (SmithKline Beecham Biologicals) describes MuV FxHN and HNxF hybrids which can be expressed in vaccinia, a mammalian cell (such as CHO) or a bacterial cell. Examples B) 3 and 4 describe the expression of  $s^+FHNa^+xFa^+$  and  
25  $Fs^+a^+xHN^+$  in CHO cells however no details are given describing the extent of expression and secretion.

Although this cited art may suggest that homochimeric paramyxoviridae glycoproteins can be expressed in a variety of cell lines including CHO cells it has  
30 now been discovered that in fact expression and secretion from CHO cells is not always successful and success cannot be predicted. Thus it has now been demonstrated that although a RSV F x G hybrid could be successfully expressed and

secreted in CHO cells, analogous homochimeric hybrids from PIV3 and MuV could not in fact be expressed in CHO cells in such manner that they could be purified from the supernatant in significant quantities.

- 5 Surprisingly, it has now been discovered that heterochimeric hybrids can be successfully expressed and secreted in both CHO and insect cells.

Accordingly in a first aspect the present invention provides a process for preparing a heterochimeric protein or an immunogenic derivative thereof comprising an  
10 immunogenic fragment of the fusion (F) protein of RSV, PIV1, PIV2, PIV3, MV or MuV and an immunogenic fragment of the attachment (G, HN or H) protein of RSV, PIV1, PIV2, PIV3, MuV or MV which process comprises expressing recombinant DNA encoding the heterochimeric protein or immunogenic derivative thereof in CHO cells and recovering the protein.

- 15 By heterochimeric protein is meant one that does not contain a fusion or attachment protein from the same pathogen.

This invention also provides novel heterochimeric proteins not previously described  
20 in WO 9314207 which can be prepared using the process of the present invention.

Thus, in a second aspect the present invention provides a heterochimeric protein or an immunogenic derivative thereof comprising an immunogenic fragment of the fusion (F) protein of RSV, PIV1, PIV2, PIV3, MV or MuV and an immunogenic  
25 fragment of the attachment (G, HN or H) protein of RSV, PIV1, PIV2, PIV3, MuV or MV, with the proviso that where one of the immunogenic fragments is derived from RSV F, RSV HN or PIV3 F, PIV3 HN, the other of the immunogenic fragments is derived from MuV F, MuV HN, MV F, MV H, PIV1 F, PIV1 HN, PIV2 F or PIV2 HN.

- 30 By an immunogenic fragment of the fusion (F) protein of RSV, PIV1, PIV2, PIV3, MV or MuV is meant a part of the protein which contains at least one antigenic

determinant capable of raising an immune response specific to the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV respectively. Included within this definition is the full length F protein, preferably however the immunogenic fragment is lacking the membrane anchor domain at its C-terminal end.

5

By an immunogenic fragment of the attachment protein (G, HN or H) of RSV, PIV1, PIV2, PIV3, MuV or MV is meant a part of the protein which contains at least one antigenic determinant capable of raising an immune response specific to the G protein of RSV, to the HN protein of PIV1, PIV2, PIV3, MuV or the H protein of MV respectively. Included within this definition is the full length G or HN protein, preferably however the immunogenic fragment is lacking the signal/anchor domain at its N-terminal end.

10

Preferably the heterochimeric protein is linked *via* an amino acid in the C-terminal part of the immunogenic fragment of the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV to an amino acid in the N-terminal part of the immunogenic fragment of the G protein of RSV, the HN protein of PIV1, PIV2, PIV3, MuV or the H protein of MV.

15

Suitably the heterochimeric protein commences at its N-terminal end with a signal sequence from the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV. Conveniently this will be part of the corresponding immunogenic fragment of the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV when this fragment is linked *via* its C-terminal end to the N-terminal end of the immunogenic fragment of the G protein of RSV, the HN protein of PIV1, PIV2, PIV3, MuV or the H protein of MV.

20

25

Alternative signal sequences may also be employed. For example, the heterochimeric protein suitably commences at its N-terminal end with a signal sequence of tissue plasminogen activator (TPA).

30

In order to enhance the level of expression the heterochimeric protein may further comprise a ubiquitin leader sequence which is suitably positioned after any signal sequence as hereinbefore described. Preferably the ubiquitin leader sequence is linked to the C-terminal end of the signal sequence of TPA.

5

Preferably the ubiquitin leader sequence is derived from yeast, for example as described in Ecker et al, J.Biological Chemistry, 1988, 264(13), 7715-7719.

10

Suitably a cleavage site is positioned between the C-terminal end of the ubiquitin sequence and the N-terminal end of the immunogenic fragment of the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV.

15

In order to facilitate chromatographic purification the heterochimeric protein suitably comprises a polyhistidine tail, for example as described in Hochuli et al, BIO/TECHNOLOGY, 1988, 1321-1325. The polyhistidine tail preferably comprises from 2 to 6 adjacent histidine residues which is suitably attached at the C-terminal end of the heterochimeric protein. Preferably a cleavage site is positioned between the polyhistidine tail and the C-terminal end of the immunogenic fragment of the G protein of RSV, the HN protein of PIV1, PIV2, PIV3, MuV or the H

20

protein of MV.

25

The cleavage site for the ubiquitin sequence and/or the polyhistidine tail may be chemical or enzymatic and preferably is an enterokinase cleavage site, for example as described in LaVallie et al, BIO-TECHNOLOGY, 1993, 187-193.

30

Following expression and purification, treatment with an enterokinase will cleave off any ubiquitin and/or polyhistidine sequence releasing the desired heterochimeric protein.

Particular heterochimeric proteins of this invention include:

the F protein of RSV lacking its membrane domain linked at its C-terminal end to the HN protein of MuV lacking its signal/anchor domain herein referred to as:

Fs<sup>+</sup>a<sup>-</sup>RSVxHNs<sup>-</sup>a<sup>-</sup>MuV, as well as  
Fs<sup>+</sup>a<sup>-</sup>PIV3 x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup> MuV;  
Fs<sup>+</sup>a<sup>-</sup> MuV x Gs<sup>-</sup>a<sup>-</sup>RSV; and  
Fs<sup>+</sup>a<sup>-</sup> MuV x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup>PIV3, and  
5 immunogenic derivatives thereof.

The present invention also provides particular heterochimeric proteins which include:

10 Fs<sup>+</sup>a<sup>-</sup>MuVxHs<sup>-</sup>a<sup>-</sup>MV; or  
Fs<sup>+</sup>a<sup>-</sup>RSVxHNs<sup>-</sup>a<sup>-</sup>PIV1; or  
Fs<sup>+</sup>a<sup>-</sup>RSVxHNs<sup>-</sup>a<sup>-</sup>PIV2, and  
immunogenic derivatives thereof.

15 The present invention also provides heterochimeric proteins comprising RSV and PIV3 proteins not specifically disclosed in WO9314207, which advantageously can be expressed from CHO cells.

These are:

20 Fs<sup>+</sup>a<sup>-</sup> (1-526) RSV x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup> (70-572) PIV3;  
Fs<sup>+</sup>a<sup>-</sup> (1-492) PIV3 x Gs<sup>-</sup>a<sup>-</sup> (69-298) RSV;  
Fs<sup>+</sup>a<sup>-</sup> (1-526) RSV x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup> (70-572) PIV3 bis;  
Fs<sup>+</sup>a<sup>-</sup> (1-526) RSV x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup> (70-572) PIV3 ent his, and  
sTPA (1-21) UB (1-74) ent Fs<sup>-</sup>a<sup>-</sup> (24-526) x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup> (70-572) PIV3, and  
immunogenic derivatives thereof.

25 The heterochimeric proteins of the present invention are immunogenic. The term immunogenic derivative as used herein encompasses any molecule which is a heterochimeric polypeptide which is immunologically reactive with antibodies raised to the heterochimeric protein of the present invention or parts thereof or with antibodies recognising the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV, the  
30 G protein of RSV, the HN protein of PIV1, PIV2, PIV3, MuV, the H protein of MV, the RSV virus, the PIV1 virus, the PIV2 virus, the PIV3 virus, the MV virus or the MuV virus, or which, when administered to a human, elicits antibodies

recognising the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV, the G protein of RSV, the HN protein of PIV1, PIV2, PIV3, MuV, the H protein of MV, the RSV virus, the PIV1 virus, the PIV2 virus, the PIV3 virus, the MV virus or the MuV virus. In particular immunogenic derivatives which are slightly longer or shorter than the heterochimeric proteins of the present invention may be used. Such derivatives may, for example, be prepared by substitution, addition, or rearrangement of amino acids or by chemical modifications thereof including the coupling or for enabling the coupling of the heterochimeric proteins to other carrier proteins such as tetanus toxoid or Hepatitis B surface antigen. All such substitutions and modifications are generally well known to those skilled in the art of peptide chemistry.

Immunogenic fragments of the heterochimeric proteins which may be useful in the preparation of vaccines may be prepared by expression of the appropriate gene fragments or by peptide synthesis, for example using the Merrifield synthesis (The Peptides, Vol 2., Academic Press, New York, p3).

In a further aspect of the invention there is provided recombinant DNA encoding the heterochimeric protein of the invention. The recombinant DNA of the invention may form part of a vector, for example a plasmid, especially an expression plasmid from which the heterochimeric protein may be expressed. Such vectors also form part of the invention, as do host cells into which the vectors have been introduced.

In order to construct the DNA encoding a heterochimeric protein according to the invention, cDNA containing the coding sequences of the RSV, PIV1, PIV2, PIV3, MV or MuV fusion and attachment proteins and optionally of the ubiquitin, polyhistidine and enterokinase cleavage sites may be manipulated using standard techniques [see for example Maniatis T. et al Molecular Cloning, Cold Spring Harbor Laboratory, Cold Spring Harbor N.Y. (1982)] as further described hereinbelow.

In another aspect of the invention there is described a process of enhancing the protein expression in mammalian cells by optimization of the codon usage of the nucleic acids transfected therein. Optimization of the codon usage involves the replacement of at least one non-preferred or less preferred codon in a natural gene encoding a heterochimeric protein by a preferred codon encoding the same amino acid. Highly mammalian-expressed genes have C or G at their degenerative position (third base in the codon) whereas the RSV or PIV3-prevalent codons have A or T. At least one codon, and more preferably all the codons of the RSV or PIV3 protein can be changed to fit at best the human usage, that is, the one (or ones) that is the most prevalent as shown below.

Ala: GCC	Cys: TGC	His: CAC	Met: ATG	Thr: ACC
Arg: CGC AGG CGG	Gln: CAG	Ile: ATC	Phe: TTC	Trp: TGG
Asn: AAC	Glu: GAG	Leu: CTG	Pro: CCC	Tyr: TAC
Asp: GAC	Gly: GGC	Lys: AAG	Ser: AGC TCC	Val: GTG

Each amino acid encoded by one of these codons are then considered humanised. The ratio between the number of humanised codons versus the total number of amino acids gives a percentage of humanisation as shown below.

- 1)  $F_{\text{RSV (1-526)original}} = 140/526 = 27\%$
- 2)  $F_{\text{RSV (1-423)humanised}} + (424-526)original = 403/526 = 77\%$
- 3)  $F_{\text{RSV (1-526)humanised}} = 489/526 = 93\%$
- 4)  $F_{\text{RSV (1-526)original}} + \text{HN}_{\text{PIV3 (70-572) original}} = 258/1029 = 25\%$
- 5)  $F_{\text{RSV (1-526)humanised}} + \text{HN}_{\text{PIV3 (70-572) original}} = 528/1029 = 51\%$
- 6)  $F_{\text{RSV (1-526)humanised}} + \text{HN}_{\text{PIV3 (70-572) humanised}} = 96\%$

The invention also provides DNA encoding a heterochimeric protein or immunogenic derivative thereof in which the codon usage of one or more nucleic acids has been substantially optimised and a process for expressing said DNA in a CHO or insect cell.

5

There have been a number of reports that have described a substantial amelioration of protein expression in mammalian cells after re-engineering the nucleic acid sequence of the heterologous protein to fit the codon usage found in highly expressed human genes (Haas J., Park E-C. and Seed B., Codon usage limitation in the expression of HiV-1 envelope glycoprotein, Current Biology, 1996, 6, n°3, 315-325 ; Kim C. H., Oh Y. and Lee T.H., Codon optimization for high-level expression of human erythropoietin (EPO) in mammalian cells, Gene 199, 1997, 293-301 ; Zolotukhin S., Potter M. Hauswirth W.W. Guy J. and Muzyczka N. A Humanized green fluorescent protein cDNA adapted for high level expression in mammalian cells. J. of Virology, July 1996, 70, n°7, 4646-4654).

10

15

Vectors comprising such DNA, hosts transformed thereby and the truncated or hybrid proteins themselves, expressed as described hereinbelow all form part of the invention.

20

For expression of the proteins of the invention, plasmids may be constructed which are suitable either for transfer into vaccinia virus or transfection into CHO cells, insect cells or Vero cells. Suitable expression vectors are described hereinbelow. Preferably the proteins of the present invention are expressed in CHO or insect cells.

25

For expression in vaccinia a vaccinia transfer plasmid such as pULB 5213 which is a derivative of pSC11 (Chakrabati *et al*, Molecular and Cellular Biology 5, 3403 - 3409, 1985) may be used. In one aspect the protein may be expressed under the control of the vaccinia P7.5 promoter.

30

For expression in CHO-K1 cells a glutamine synthetase (GS) vector such as pEE14 may suitably be used so that the protein is expressed under the control of the major immediate early promoter of human cytomegalovirus (hCMV-MIE). Alternatively a vector which allows the expression of the coding module as a polycistronic transcript with the *neo* selection gene may suitably be used. In one preferred aspect the coding module is under the control of the Rous Sarcoma Long Terminal Repeat (LTR) promoter.

Preferably the plasmid for expression in CHO-K1 cells carries a GS expression cassette suitable for gene amplification using methionine sulfoximine (MSX). Alternatively the plasmid for expression in CHO-K1 cells carries a DHFR expression cassette suitable for gene amplification using methotrexate (MTX).

Preferably expression of the heterochimeric protein of the present invention is carried out in the presence of sodium butyrate and/or dimethyl sulfoxide (DMSO) which may enhance gene expression.

For expression in insect cells a shuttle vector such as pAcUW51 or pAcGP67 may be used. In one aspect the protein may be expressed under the control of the baculovirus p10 promoter or the polyhedrin promoter.

The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and *in vivo* infection with this live vector will lead to *in vivo* expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g. vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), *Listeria*, *Salmonella*, *Shigella*, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

In yet another aspect of the invention there is provided a vaccine composition comprising a heterochimeric protein or immunogenic derivative thereof according to the invention in combination with a pharmaceutically acceptable carrier, a protein according to the invention for use in vaccinating a mammal and the use of a protein according to the invention in the preparation of a vaccine.

Optionally, and advantageously, the vaccine of the present invention is combined with other immunogens to afford a polyvalent vaccine. In a preferred embodiment the heterochimeric protein is combined with other subcomponents of RSV, PIV1, PIV2, PIV3, MuV or MV, e.g. the single proteins F, G, HN or H or homochimeric proteins such as RSV FxG, PIV3 FxHN or MuV FxHN.

In a particular aspect the invention further provides a vaccine composition comprising a protein according to the invention together with a suitable carrier or adjuvant.

Vaccine preparation is generally described in *New Trends and Developments in Vaccines*, edited by Voller *et al.*, University Park Press, Baltimore, Maryland, U.S.A., 1978. Encapsulation within liposomes is described, for example by Fullerton, U.S. Patent 4,235,877.

In the vaccine of the present invention, an aqueous solution of the protein(s) can be used directly. Alternatively, the protein, with or without prior lyophilisation, can be mixed, absorbed or adsorbed with any of the various known adjuvants. Such adjuvants include, but are not limited to, aluminium hydroxide, muramyl dipeptide and saponins such as Quil A. Particularly preferred adjuvants are MPL (monophosphoryl lipid A) and 3D-MPL (3 deacylated monophosphoryl lipid A) [US patent 4,912,094], optionally formulated with aluminium hydroxide (EP 0 689 454) or oil in water emulsions (WO 95/17210). A further preferred adjuvant is known as QS21 which can be obtained by the method disclosed in US patent 5,057,540. Use of 3D-MPL is described by Ribí *et al.* in *Microbiology* (1986) Levie *et al.* (eds)

Amer. Soc. Microbiol. Wash. D.C., 9-13. Use of Quil A is disclosed by Dalsgaard *et al.*, (1977), Acta Vet Scand, 18, 349. Use of combined 3D-MPL and QS21 is described in WO 94/00153 (SmithKline Beecham Biologicals s.a). QS21 may be advantageously formulated with cholesterol containing liposomes, wherein 3D-MPL  
5 is present either in solution or incorporated in the membrane, as described in WO 96/33739.

As a further exemplary alternative, a heterochimeric protein of the invention or an immunogenic fragment thereof can be encapsulated within microparticles such as  
10 liposomes or associated with oil-in-water emulsions. Encapsulation within liposomes is described by Fullerton in US patent 4,235,877. In yet another exemplary alternative, a heterochimeric protein according to the invention or an immunogenic fragment thereof can be conjugated to an immunostimulating macromolecule, such as killed *Bordetella* or a tetanus toxoid. Conjugation of  
15 proteins to macromolecules is disclosed, for example by Likhite in patent 4,372,945 and Armor *et al.* in US patent 4,474,757.

The amount of the protein of the present invention present in each vaccine dose is selected as an amount which induces an immunoprotective response without  
20 significant, adverse side effects in typical vaccines. Such amount will vary depending upon which specific immunogen is employed and whether or not the vaccine is adjuvanted. Generally, it is expected that each dose will comprise 1-1000µg of protein, preferably 1-200 µg. An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of antibody  
25 titres and other responses in subjects.

The following examples and the attached figures (explained below) illustrate the invention.

30 **In the Figures:**

**Figure 34A** shows the impact of humanisation on the level of expression of FrHNp, where:

FhHNE10 = product expressed by the pEE14FhHN transfected clone E10;

FhHNE7 = product expressed by the pEE14FhHN transfected clone E7;

FHNbis = product expressed by the pEE14FHN transfected clone;

+but = 2mM Nabutyrate has been added to the cell medium, 3 days before

5 harvest;

pEE14 = negative control;

Fdroso = purified Fa- (drosophila derived); the standard protein in this ELISA assay wherein 1ul of standard corresponds to 1ng of product.

Figure 34B shows humanisation impact on the level of expression of  $F_{RSV}HN_{PIV3}$ ,

10 where the level of expression was determined by ELISA. Fdroso = purified Fa- (drosophila derived) that is the standard protein in this ELISA assay, 1ul of standard corresponds to 1ng of product.

## EXAMPLES

### Example 1

In order to vaccinate with a single immunogen, heterochimeric DNA molecules  
5 were constructed combining extracellular domains of the F and the attachment  
protein for each virus. DNA constructs for the PIV3 and MuV have already been  
described in WO9306218 and WO9425600, respectively. The DNA molecule,  
combining the extracellular domains of the RSV F and G proteins were constructed  
as described below.

10

The DNA pieces were first inserted into the mammalian expression vector based on  
the replicon of the Semliki Forest Virus (pSFV1). This expression system does not  
lead to a stable expression mammalian cell line but, however gives an indication  
whether or not the chimeric protein is expressed and whether the product is  
15 effectively secreted in the culture medium, which is advantageous for the  
purification procedure.

Stable expression in the culture medium of mammalian cell lines is preferred to  
obtain good quality and quantities of paramyxovirus glycoproteins. All the chimeric  
20 modules have been inserted in the shuttle vector, the pEE14, which integrates in the  
genome of mammalian cells such as CHO-K1. A quite good expression level was  
obtained with the RSV FxG homochimeric recombinant protein, however negligible  
expression was obtained for the FxHN recombinant homochimeric protein of either  
PIV3 or MuV. Expression of heterochimeric proteins was obtained from CHO  
25 cells.

Thus by constructing heterochimeric DNA molecules combining the extracellular  
domains of the F protein of one virus linked to the extra cellular domain of the HN  
or G protein of another virus and inserting them into the pEE14 vector for CHO  
30 expression it has been possible to raise the expression level of these proteins. These  
proteins may be used to achieve protection against at least two paramyxoviridae  
viruses with a single immunogen.

Some of the chimeric molecules have been inserted into the shuttle vectors, pAcUW51 and pACGP67, which integrate in the genome of bacterial and lepidopteran cells. Surprisingly good expression of heterochimeric proteins was  
5 obtained from insect cells.

## Vector construction

### Preliminary Constructs

10

#### a) Plasmid pNIV2819

Starting from plasmid pNIV2801, a cDNA clone encoding *inter alia* the F protein of RSV (type RSS-2; received from Dr Pringle, UK) we reconstructed a cDNA  
15 module coding for the F protein lacking the membrane anchor sequence.

Plasmid pNIV2801 was digested with *Pst*I in order to recover a 1416 bp DNA piece encoding amino acid residues 18 to 489 of the F protein. Synthetic  
oligonucleotides, specifying respectively the sequences for amino acids 1 to 17 and  
20 490 to 526, were used to produce the corresponding cDNA fragments by the polymerase chain reaction performed with pNIV2801 DNA as template. The primers were designed to generate also unique flanking restriction sites useful for subsequent cloning steps. The coding module was assembled, by ligation, from the three DNA pieces described above and introduced into the standard cloning vector  
25 pUC19, to create plasmid pNIV2819. This plasmid encodes the RSV F protein carrying its signal sequence but lacking its anchor sequence (figure 1).

#### b) Plasmid pNIV 2820

30 The cDNA module encoding the full length F protein of RSV was constructed as follows. Using two synthetic oligonucleotides, the polymerase chain reaction was performed with pNIV2801 DNA as template to generate a 273 bp DNA fragment

encompassing the sequence coding for aa 490 to aa 574 of the F protein, the stop codon and unique restriction sites useful for subsequent cloning steps. This fragment was digested with *NsiI* and *EcoRI* and substituted for the *NsiI-EcoRI* DNA piece present in the coding module of pNIV2819 (figure 2). The resulting plasmid, pNIV2820, thus encodes the RSV F protein carrying both signal and membrane anchor sequences.

#### c) Plasmid pNIV2841

10 In this construction, the DNA coding for aa 165 to 176 of the G protein of RSV is fused to the DNA encoding the RSV  $Fs^+a^-$  protein. This part of the G protein is conserved among both subgroups of RSV.

The starting material, pNIV2819, was digested by *NcoI* and *SmaI* yielding a 1601 bp fragment. This fragment was subcloned into the *NcoI* and *MscI* sites of pNIV103 (a derivative of pULB1221, see European Patent Application No. 186643) leading to pNIV2844. This subcloning allowed to place the translation initiation site of the F protein in a more favourable context according to the model proposed by Kozak (Kozak M, Nature 308, 241-246, 1984).

20

A 1605 bp fragment was recovered from pNIV2844 by digestion with *KpnI* and *SalI* and introduced by ligation into pUC19 digested with *KpnI* and *SalI*, creating pNIV2840.

25 Two complementary synthetic oligonucleotides specifying the sequence for amino acids 165 to 176 of the G protein followed by a stop codon and flanked by *NsiI*, *BamHI*, *EcoRI* and *HindIII* sites were hybridized. The 55 bp resulting fragment was cloned into the pNIV2840 digested by *NsiI* and *HindIII*, thus replacing a 142 bp DNA sequence encoding amino acids 491 to 526 of the F protein. The resulting recombinant plasmid, pNIV2841, thus contains the sequence coding for amino acids 1 to 490 of the F protein followed by amino acids 165 to 176 of the G protein (figure 3).

30

### Vector Construction

#### I) For transfer into the pSFV1 vector

- 5 a) The RSV fusion protein lacking the membrane anchor domain fused to the MuV hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{RSV}$  (1-526)  $HN_{MuV}$  (60-582).

Plasmid pNIV2875, a derivative of pNIV2820 which carries the DNA coding for  
10 the F protein of RSV in which the *SpeI* restriction site has been eliminated by site-directed mutagenesis into the pUC19 vector, has been digested by *HindIII* and *BspHI*, and a 1618 bp fragment has been isolated. Plasmid pNIV3229, a derivative of pNIV3215 whose construction has been already described in WO9425600 and which carries the DNA coding for the HN protein of MuV into the pUC19 vector,  
15 has been digested with *BbsI* and *BamHI*; a 1580 bp fragment has been isolated. Both fragments were linked together by two complementary synthetic *BspHI-BbsI* oligonucleotides (Fig 4A) restoring the coding sequence of the chimeric molecule and were inserted into the *BamHI-HindIII* site of the pUC19 vector leading to pNIV4102. (Fig4B) After the sequencing of the junction regions, the chimeric  
20 cassette was retrieved from pNIV4102 by a *BamHI* digestion and was inserted into the *BamHI* site of the pSFV1 vector (Liljeström, P. and Garoff, H. (1991) Bio/Technology 9, 1356). The resulting plasmid, pNIV4104, contains into the pSFV1 vector the sequence coding for amino acids 1 to 526 of the RSV F protein followed by amino acids 60 to 582 of the MuV HN protein. (Fig4C)

25

- b) The RSV fusion protein lacking the membrane anchor domain fused to the PIV3 hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{RSV}$  (1-526)  $HN_{PIV3}$  (70-572).

- 30 Plasmid pIBI-HN, a cDNA clone containing the complete coding sequence of protein HN of PIV3 as well as its 3' non coding sequence (received from Dr. K. Dimock, University of Ottawa, Canada), has been digested by *AseI* and *BamHI* and

a 1468 bp fragment has been isolated. Plasmid pNIV2875 (see supra), which carries the DNA encoding the F protein of RSV, in which the unique *SpeI* site has been eliminated by site-directed mutagenesis, inserted into the pUC19 vector, has been digested by *BamHI* and *BspHI*, and a 1588 bp fragment has been isolated. Both  
5 fragments were linked together by two complementary synthetic *BspHI*-*AseI* oligonucleotides (Fig5A) and were inserted into the *BamHI* site of the pUC19 vector leading to pNIV4105 or to pNIV4109 (Fig5B) depending of the orientation of the chimeric module in the vector. After the sequencing of the junction region, the chimeric cassette was retrieved by a *BamHI* digestion from pNIV4109 and inserted  
10 into the *BamHI* site of the pSFV1 vector. The resulting plasmid, pNIV4110, contains, inserted into the pSFV1 vector, the sequence coding for amino acids 1 to 526 of the RSV F protein followed by amino acids 70 to 572 of the PIV3 HN protein. (Fig5C)

- 15 c) The PIV3 fusion protein lacking the membrane anchor domain fused to the RSV attachment protein lacking the signal-anchor domain,  $F_{PIV3}$  (1-492)  $G_{RSV}$  (69-298).

Plasmid pNIV3310, described in WO9306218 which carries the DNA coding for  
20 amino acids 1 to 484 of the PIV3 F protein followed by amino acids 87 to 572 of the PIV3 HN protein into the pIBI vector, was digested by *EcoRI* and *BgIII*, and a 1435 bp fragment has been isolated. Plasmid pNIV2850, which carries the RSV G protein into the pUC19 vector, has been digested by *MaeIII* and *HindIII*, and a 694 bp fragment has been isolated. Both fragments were then linked together by using  
25 two complementary *BgIII*-*MaeIII* synthetic linkers (Fig6A) and were inserted into the *EcoRI*-*HindIII* sites of pUC19 vector leading to pNIV4103 (Fig6B). The chimeric module was then retrieved from the pUC19 vector by a *BamHI*-*HindIII* digestion. After treating the protruding ends with the Klenow polymerase, the chimeric cassette has been inserted into the *SmaI* site of pSFV1 vector. The  
30 resulting plasmid pNIV4106, thus contains the sequence coding for amino acids 1 to 492 of the F protein of PIV3 followed by amino acids 69 to 298 of the G protein of RSV inserted into the pSFV1 vector (Fig6C).

d) The PIV3 fusion protein lacking the membrane anchor domain linked to the MuV hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{PIV3}$  (1-493)  $HN_{MuV}$  (60-582).

5

Plasmid pNIV3310 (see supra,  $FHN_{PIV3}$  in pIBI) was digested by *EcoRI* and *BglII* and a 1435 bp fragment was isolated. Plasmid pNIV3229 (see supra,  $HN_{MuV}$  into pUC19) was digested by *BbsI* and *HindIII*, and a 1610 bp fragment was isolated. Both fragments were linked together by adding two synthetic complementary linkers specifying a *BglII* and a *BbsI* ends (Fig7A) into the pUC19 vector leading to pNIV4117 (Fig7B). After sequencing the junction region, the chimeric cassette was retrieved from the pUC19 vector by a *BamHI* digestion and was inserted into the *BamHI* site of the pSFV1 vector. The resulting plasmid pNIV4118 encodes, cloned in the pSFV1 vector, the DNA sequence specifying amino acids 1 to 493 of the PIV3 fusion protein linked to amino acids 60 to 582 of the MuV HN protein (Fig7C).

e) The MuV fusion protein lacking its membrane anchor domain linked to the RSV attachment protein lacking its signal-anchor domain,  $F_{MuV}$  (1-482)  $G_{RSV}$  (69-298).

20

Plasmid pNIV3221, described in WO9425600 which carries the sequence encoding amino acids 1 to 462 of the MuV fusion protein within the pUC19 vector, has been digested with *EcoRI* and *BsrFI*, and a 771 bp fragment has been purified. Plasmid pNIV3221 has been also digested with *BsrFI* and *PstI*, and a 628 bp fragment has been isolated. Plasmid pNIV2850 (see supra,  $G_{RSV}$  into the pUC19) has been digested with *MaeIII* and *HindIII* and a 694 bp fragment has been isolated. The three fragments were linked together; the  $F_{MuV}/G_{RSV}$  junction was created by adding to the ligation reaction two synthetic complementary oligonucleotide specifying *PstI* and *MaeIII* sites (Fig8A), and were inserted into the *EcoRI-HindIII* sites of the pBluescript vector leading to pNIV4113 (Fig8B). The chimeric cassette was recovered from pNIV4113 by a *Asp718I* digestion and, after treating the protruding

30

ends with the Klenow polymerase, was inserted into the *Sma*I site of the pSFV1 vector. The resulting plasmid, pNIV4114 contains into the pSFV1 vector the sequence specifying amino acids 1 to 482 of the MuV F protein linked to amino acids 69 to 298 of the RSV G protein (Fig8C).

5

f) The MuV fusion protein lacking its membrane anchor domain linked to the PIV3 hemagglutinin-neuraminidase lacking its signal-anchor domain,  $F_{\text{MuV}}$  (1-482)  $HN_{\text{PIV3}}$  (54-572).

10 Plasmid pNIV4113 (see supra,  $F_{\text{MuV}}$  x  $G_{\text{RSV}}$  in pBluescript) was digested by *Bsa*I and *Bam*HI, a 1469 bp fragment was isolated. Plasmid pNIV3308, described in WO9306218 and which carries the DNA sequence specifying amino acids 1 to 31 followed by amino acids 54 to 572 of the PIV3 HN protein into the pIBI vector, was digested by *Eco*RI and *Bam*HI and a 1569 bp fragment was isolated. Both  
15 fragments were linked together by two synthetic complementary linkers specifying *Bsa*I and *Eco*RI sites (Fig9A) into the *Bam*HI site of pBluescript leading to pNIV4115 (Fig9B). The chimeric module was recovered from pNIV4115 by a *Bam*HI digestion and was inserted into *Bam*HI site of pSFV1 vector. The resulting plasmid, pNIV4116, encodes, in the pSFV1 vector, the sequence specifying amino  
20 acids 1-482 of the MuV F protein fused to amino acids 54 to 572 of the PIV3 HN protein (Fig9C).

g) The RSV fusion protein lacking its membrane anchor domain linked to the RSV attachment protein lacking its signal-anchor domain,  $F_{\text{RSV}}$  (1-526)  $G_{\text{RSV}}$  (69-  
25 298).

Plasmid pNIV2857 (Fig16A), a derivative of pNIV2841 and which contains the DNA sequence coding for amino acids 1 to 526 of the RSV fusion protein linked to amino acids 69 to 298 of the RSV attachment protein, has been digested by *Asp*718I  
30 and *Hind*III and a 2180 bp fragment has been isolated. After treating the protruding extremities with Klenow's polymerase, this fragment has been inserted in the *Sma*I site of the pSFV1 vector. The resulting plasmid pNIV2870, contains in the pSFV1

vector, the DNA sequence coding for amino acids 1 to 526 of the RSV fusion protein linked to amino acids 69 to 298 of the RSV attachment protein (Fig16B).

## II) For transfection into CHO cells

5

a) The RSV fusion protein lacking the membrane anchor domain fused to the MuV hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{RSV}$  (1-526)  $HN_{MuV}$  (60-582).

10 Plasmid pNIV4102, (Fig10A, see supra,  $F_{RSV}$  x  $HN_{MuV}$  into the pUC19 vector) has been digested with *Bam*HI, and after treating the protruding ends with the Klenow polymerase, the chimeric module has been inserted into the *Sma*I site of the glutamine synthetase (GS) vector, pEE14 (Cockett *et al.*, 1990, Bio/Technology 8, 662-667). The resulting plasmid pEE14  $Fs^+a^-$  RSV x  $HNs^+a^-$  MuV contains  
15 sequences coding for amino acids 1 to 526 of the RSV F protein fused to amino acids 60 to 582 of the MuV HN protein under the control of the major immediate early promoter of the human cytomegalovirus (hCMV-MIE) (Fig10B).

b) The RSV fusion protein lacking its membrane anchor domain linked to the  
20 PIV3 hemagglutinin-neuraminidase lacking its signal-anchor domain,  $F_{RSV}$  (1-526)  $HN_{PIV3}$  (70-572).

Plasmids pNIV4105 and pNIV4109 (Fig11A and B, see supra,  $F_{RSV}$  x  $HN_{PIV3}$  into the pUC19 vector) were digested by *Eco*RI and *Xho*I and a 2032 bp as well as a  
25 1064 bp fragments were isolated. Both fragments were inserted together into the *Eco*RI site of pEE14. The resulting plasmid pEE14  $Fs^+a^-$  RSV x  $HNs^+a^-$  PIV3 contains sequences coding for amino acids 1 to 526 of the RSV F protein fused to amino acids 70 to 572 of the PIV3 HN protein under the control of the hCMV promoter (Fig11C).

30

c) The PIV3 fusion protein lacking the membrane anchor region linked to the RSV attachment protein lacking the signal-anchor domain,  $F_{PIV3}$  (1-492)  $G_{RSV}$  (69-298).

5 Plasmid pNIV4103 (Fig12A, see supra,  $F_{PIV3}$  x  $G_{RSV}$  into the pUC19 vector) was digested by *HindIII* and a 2180 bp fragment was isolated. After treating the protruding extremities with the Klenow polymerase, the chimeric module was inserted into the *SmaI* site of the pEE14 vector. The resulting plasmid, pEE14  $Fs^+a^-$  PIV3 x  $Gs^-a^-$  RSV, contains, under the control of the hCMV promoter, the sequence  
10 encoding amino acids 1 to 492 of the PIV3 F protein followed by amino acids 69 to 298 of the RSV G protein (Fig 12B).

d) The PIV3 fusion protein lacking the membrane anchor domain fused to the MuV hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{PIV3}$  (1-493)  $HN_{MuV}$  (60-582).  
15

Plasmid pNIV4117 (Fig13A, see supra,  $F_{PIV3}$   $HN_{MuV}$  into the pUC19 vector) was digested with *HindIII* and a 3119 bp fragment was isolated and inserted into the *HindIII* site of the pEE14 vector. The resulting plasmid, pEE14  $Fs^+a^-$  PIV3 x  $HNs^-a^-$  MuV, contains under the control of the hCMV promoter a sequence encoding  
20 amino acids 1 to 493 of the PIV3 fusion protein fused to amino acids 60 to 582 of the MuV HN protein (Fig13B).

e) The MuV fusion protein lacking its membrane anchor domain fused to the RSV attachment protein lacking its signal-anchor domain,  $F_{MuV}$  (1-482)  $G_{RSV}$  (69-298).  
25

Plasmid pNIV4113 (Fig14A, see supra,  $F_{MuV}$   $G_{RSV}$  into the pBluescript vector) has been digested *Asp718I*, the protruding ends have been treated by the Klenow  
30 polymerase. A 2200 bp fragment has been isolated and inserted into the *SmaI* site of pEE14. The resulting plasmid, pEE14  $Fs^+a^-$  MuV x  $Gs^-a^-$  RSV, has, under the

control of the hCMV promoter, the sequence encoding amino acids 1 to 482 of the MuV F protein followed by amino acids 69 to 298 of the RSV G protein (Fig14B).

- f) The MuV fusion protein lacking its membrane anchor domain fused to the PIV3 hemagglutinin-neuraminidase lacking its signal-anchor domain,  $F_{\text{MuV}}$  (1-482)  $\text{HN}_{\text{PIV3}}$  (54-572).

Plasmid pNIV4115 (Fig15A, see supra,  $F_{\text{MuV}}$  x  $\text{HN}_{\text{PIV3}}$  into the pBluescript vector) has been digested with *EcoRI* and a 3040 bp fragment has been inserted into the *EcoRI* site of the pEE14 vector. The resulting plasmid, pEE14  $F_{\text{MuV}}$  x  $\text{HN}_{\text{PIV3}}$ , contains, downstream to the hCMV promoter region, a sequence coding for amino acids 1 to 482 of the MuV F protein followed by amino acids 54 to 572 of the PIV3 HN protein (Fig15B).

- g) The RSV fusion protein lacking its membrane anchor domain linked to the RSV attachment protein lacking its signal-anchor domain,  $F_{\text{RSV}}$  (1-526)  $G_{\text{RSV}}$  (69-298).

Plasmid pNIV2857 (Fig17A), a derivative of pNIV2841 and which contains the DNA sequence coding for amino acids 1 to 526 of the RSV fusion protein linked to amino acids 69 to 298 of the RSV attachment protein, has been digested by *Asp718I* and *HindIII* and a 2180 bp fragment has been isolated. After treating the protruding extremities with Klenow's polymerase, this fragment has been inserted the *SmaI* site of the pEE14vector. The resulting plasmid, pEE14  $F_{\text{RSV}}$  x  $G_{\text{RSV}}$ , contains under the control of the hCMV promoter the DNA sequence coding for amino acids 1 to 526 of the RSV fusion protein linked to amino acids 69 to 298 of the RSV attachment protein (Fig17B).

- h) The original RSV fusion protein lacking the membrane anchor domain linked to the PIV3 hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{\text{RSV}}$  (1-526)  $\text{HN}_{\text{PIV3}}$  (70-572) bis.

Plasmid pNIV2852, a derivative of pNIV2820 which carries the DNA encoding the RSV F protein where the translation initiation site is in a more favourable context according to the model proposed by Kozak (Kozak M, Nature 308, 241-246, 1984), has been digested *BamHI* and *BspHI*, and a 1588 bp fragment has been isolated.

5

Plasmid pIBI-HN, a cDNA clone containing the complete coding sequence of the HN protein of PIV3 (received from Dr. K. Dimock, University of Ottawa, Canada) has been digested by *AseI* and *BamHI* and a 1468 bp has been isolated.

- 10 Both fragments were linked together by two complementary synthetic *BspHI-AseI* adaptators (Fig18A) and were inserted into the *BamHI* site of the pUC19 vector leading to pNIV4120 (Fig18B).

- 15 After the sequencing of the junction region, the chimeric cassette was retrieved by a *BamHI* digestion from pNIV4120 and inserted into the *BamHI* compatible *BclI* site of the pEE14 vector. The resulting plasmid pEE14  $F_{RSV}^+ a_{RSV} \times HNs a_{PIV3}$  bis contains the sequences coding for amino acids 1 to 526 of the RSV F protein fused to amino acids 70 to 572 of the PIV3 HN protein under the control of the hCMV promoter (Fig18C).

20

- This construct differs from the earlier pEE14  $F_{RSV}^+ a_{RSV} \times HNs a_{PIV3}$  construct (II-a) in the F coding region. In  $F_{RSV}HN_{PIV3}$  bis, the nucleic acid sequence found in  $F_{RSV}HN_{PIV3}$ , ATG GAT CTG (those codons are specifying aa Met1, Asp2 and Leu3) and ACC AGT (specifying aa Thr54 and Ser 55) is replaced by the original  
25 sequence of the RSV F protein that is ATG GAG TTG (specifying aa Met1, Glu2, Leu3) and ACT AGT (specifying Thr54 and Ser55).

- i) The original RSV fusion protein lacking the membrane anchor domain linked to the PIV3 hemagglutinin-neuraminidase lacking the signal-anchor domain  
30 with, at the C-terminal part, a polyhistidine tail preceded by the enterokinase cleavage site,  $F_{RSV}$  (1-526)  $HN_{PIV3}$  (70-572) en his

Plasmid pIBI-HN, a cDNA containing the PIV3 HN protein coding sequence (see supra) has been digested by *PstI* and *SphI*. A 4588 bp fragment has been isolated and linked to complementary synthetic *PstI-SphI* adaptators (Fig19A).

- 5 After the sequencing of the junctions as well as the synthetic linkers, the resulting plasmid pNIV3340 has been digested by *XhoI* and *BamHI* and a 1121 bp fragment has been isolated (Fig19B).

- 10 Plasmid pNIV4120 (see supra) has been digested by *XhoI* and *BamHI* and a 2017 bp fragment has been isolated (Fig19C).

- Both fragments were linked together and inserted into the *BamHI* compatible *BclI* site of the pEE14 vector. The resulting plasmid pEE14 FRSVs<sup>+</sup> x HN<sup>a</sup> en his contains, under the control of the hCMV promoter, sequences coding for amino acids 1 to 526 of the RSV fusion protein fused to the amino acids 70-572 of the PIV3 HN protein fused to the enterokinase cleavage site, ({Asp} x4 Lys) followed by a polyhistidine tail ({his}x6) and a stop codon (Fig19D).

- 20 j) The signal domain of the tissue plasminogen activator fused to the yeast ubiquitin followed by the enterokinase cleavage recognition site and the original RSV fusion protein lacking its membrane signal and anchor domains linked to the PIV3 hemagglutinin-neuraminidase lacking the signal-anchor domain, sTPA(1-21) UB(1-74) ent F<sub>RSV</sub> (24-526) HN<sub>PIV3</sub> (70-572)bis.

25

- 1) The signal domain of the tissue plasminogen activator fused to the yeast ubiquitin.

- 30 A 208 bp fragment corresponding to amino acid 1 to 76 of the ubiquitin protein of *Saccharomyces cerevisiae* was isolated by a digestion of pNIV3475 ( a derivative of YEPUBSTUALL, a yeast 2  $\mu$  vector backbone carrying the yeast ubiquitin) with *BamHI* and *XbaI* (Fig 20A).

Plasmid JW4304 (received from J. Mullins, University of Washington, U.S.A) which encodes the signal domain of the tissue plasminogen activator (sTPA) was digested by *NheI* and *BamHI* and a 5115bp was isolated. Both fragments were  
5 linked together using two synthetic complementary *NheI-XbaI* adaptators (Fig20B). The resulting plasmid pNIV4121 was digested by *HindIII* and *BamHI*. A 330 bp fragment was isolated and inserted into the *HindIII* and *BamHI* sites of the pBluescript vector. The resulting plasmid pNIV4122 contains the DNA sequence specifying the signal domain of the tissue plasminogen activator followed by an  
10 alanine and a serine residue (those two amino acids are known to produce a good leader cleavage) fused to the yeast ubiquitin (Fig 20C).

2) The signal domain of the tissue plasminogen activator linked to the yeast ubiquitin followed by the enterokinase cleavage recognition site and amino acid  
15 24 to 55 of the original fusion protein of RSV.

Plasmid pNIV4122 (Fig 21A, see supra) was digested by *AflIII* and *SpeI*. A 3212 bp fragment was isolated and linked to synthetic complementary *AflIII-SpeI* adaptators (Fig21B). The entire module was then sequenced. The resulting plasmid pNIV4123  
20 encodes the signal domain of the tissue plasminogen activator linked to the N-terminal 74 aa of the yeast ubiquitin followed by the recognition site of enterokinase {(Asp)<sup>4</sup> Lys} and amino acid 24 to 55 of the original fusion protein of RSV (Fig21C).

25 3) The signal domain of the tissue plasminogen activator linked to the yeast ubiquitin followed by the enterokinase cleavage recognition site and the RSV fusion protein linked to the PIV3 hemagglutinin-neuraminidase lacking their membrane domains.

30 Plasmid pNIV4123 (Fig 22A, see supra) was digested by *HindIII*, treated by the Klenow polymerase and digested by *SpeI*. A 408 bp fragment has been isolated.

Plasmid pNIV4120 (Fig 22B, see supra) has been digested by *XbaI*, treated by the Klenow polymerase, and digested by *SpeI*. A 5620 bp fragment has been isolated.

Both fragment have been linked together to generate pNIV4124 (Fig 22C).

5

The entire coding module was retrieved from pNIV4124 by a digestion with *XbaI* and *EcoRI* and was inserted into the *XbaI* and *EcoRI* sites of the pEE14 expression vector. The resulting plasmid pEE14 sTPA x UBI x EN x Fs a RSV x HN s a PIV3, contains, under the control of the hCMV promoter, the sequence coding for aa1-21 of the tissue plasminogen activator followed by an alanine and a serine residue, by the 74 N-terminal amino acids of the yeast ubiquitin, by the recognition cleavage site of the enterokinase ({Asp}4 Lys), by aa 24-526 of the original RSV fusion protein and by aa 70-572 of the hemagglutinin-neuraminidase of PIV3.

10

### 15 III) For transfection into Insect Cells

a) The original RSV fusion protein lacking the membrane anchor domain linked to the PIV3 hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{RSV}$  (1-526)  $HN_{PIV3}$  (70-572) bis.

20

Plasmid pNIV4120 (FIG 23A) was digested by *BamHI* and a 3114 bp fragment was isolated and inserted into the *BamHI* site of the baculovirus transfer vector, pAcUW51 (PharMingen). The resulting plasmid pNIV4132 (Fig 23B) contains, under the control of the polyhedrin promoter, the sequence coding for amino acids 1-526 of the RSV F protein fused to amino acids 70-572 of the PIV3 HN protein.

25

b) The baculovirus gp67 signal peptide fused to the original RSV fusion protein lacking both membrane signal and anchor domain linked to the PIV3 hemagglutinin-neuraminidase lacking the signal-anchor domain, sGP67 $F_{RSV}$  (25-526)  $HN_{PIV3}$  (70-572) bis.

30

Plasmid pNIV4120 (FIG 24A, see supra) was digested by *Bam*HI and *Spe*I and a 2939 bp fragment was isolated, linked to two complementary synthetic *Bam*HI-*Spe*I adaptators and inserted into the *Bam*HI site of the baculovirus transfer vector, pAcGP67A (PharMingen). The resulting plasmid pNIV4136 (Fig 24) contains,  
5 under the control of the polyhedrin promoter, the sequence coding for amino acids 1-38 of the Baculovirus gp67 protein, followed by an Alanine and an Aspartate linked to amino acids 25-526 of the RSV F protein fused to amino acids 70-572 of the PiV3 HN protein.

## 10 Expression in eukaryotic cells

### A) via the pSFV1 vector

The pSFV1 vector is based on the Semliki Forest Virus (SFV) replicon. The DNA  
15 of interest is cloned into the pSFV1 vector that serves as a template for *in vitro* synthesis of recombinant RNA. The RNA is transfected into mammalian cells such as BHK-21 cells. The recombinant RNA in the cells drives its own replication and capping resulting in production of heterologous protein.

20 Plasmids pNIV2870 was digested with *Pvu*II; pNIV4106, pNIV4110, pNIV4114, pNIV4116 and pNIV4118 were digested with *Spe*I prior to RNA transcription. After a phenol extraction followed by an ethanol precipitation, 2  $\mu$ g of linearized DNA was used as a template for RNA production. About 5  $\mu$ g RNA was used to  
25 transfect, by electroporation, about  $8 \times 10^6$  BHK-21 cells. All experimental procedures for RNA production and cell transfection are detailed in Liljestrom and Garoff (Bio/Technology, 1991, 9, 1356).

After 24 h to 48 h post-electroporation, cells and spent culture medium have been collected for ELISA and radioimmunoprecipitation assays.

30 a) pNIV4104, F<sub>RSV</sub> HN<sub>MuV</sub>

ELISA were done using mAb 2072 anti-HN MuV (Örvell, 1984, *J. Immunology* 132, 2622-2629) or 20RG45, a goat anti-RSV serum (Fitzgerald, U.S.A.) to coat the microtiter plates and a rabbit polyclonal anti-SBL-1 (MuV) serum or mAb19 anti-F RSV (G.Taylor, Inst. of Animal Health, Compton Lab., U.K.) as capture antibody.

Radioimmunoprecipitation of the  $^{35}\text{S}$ -methionine labelled product was done using mAb2072 (Örvell) and products were resolved onto 7.5% SDS-PAGE.

10 b) pNIV4110,  $\text{F}_{\text{RSV}} \text{HN}_{\text{PIV3}}$

ELISA were done using anti-RSV goat serum 20RG45 or mAb anti- $\text{HN}_{\text{PIV3}}$  4830 (Rydbeck *et al*, *J. Gen. Virol.* 67, 1531-1542, 1986) to coat microtiter plates and mAb19 anti-F RSV (G.Taylor) or rabbit anti-PIV3 (E.Norrby, Stockholm) serum as a capture antibody.

Radioimmunoprecipitation was done using anti-HN PIV3 mAb4830.

20 c) pNIV4106,  $\text{F}_{\text{PIV3}} \text{G}_{\text{RSV}}$

ELISA were done using mAb anti- $\text{F}_{\text{PIV3}}$  4549 (E.Norrby, Stockholm) or mAb anti  $\text{G}_{\text{RSV}}$  858-2 (Chemicon, U.S.A.) to coat microtiter plates and a rabbit anti-PIV3 serum as a capture antibody.

25 Radioimmunoprecipitation was done using mAb anti- $\text{F}_{\text{PIV3}}$  3283 (Behringwerke).

d) pNIV4118,  $\text{F}_{\text{PIV3}} \text{HN}_{\text{MuV}}$

ELISA plates were coated with anti-F PIV3 mAb 1031215 (Norrby) or with mAb 2072 anti-HN MuV (Örvel) and rabbit anti-PIV3 sera or rabbit anti-MuV sera were used as capture antibody.

Immunoprecipitation of labelled product was done using mAb 2072 anti-HN MuV.

e) pNIV4114,  $F_{\text{MuV}} \times G_{\text{RSV}}$

ELISA plates were coated with anti-F MuV monoclonal 5414 (Örvell) or anti  $G_{\text{RSV}}$   
5 mAb (Chemicon) and a rabbit anti-SBL-1 serum was used as a capture antibody.

f) pNIV4116,  $F_{\text{MuV}} \times \text{HN}_{\text{PIV3}}$

ELISA plates were coated with anti-F MuV mAb 5414 (Örvell) or mAb anti-HN  
10 PIV3 4830 (Norrby) and rabbit anti-SBL-1 serum or a rabbit anti-PIV3 serum as a  
capture antibody.

g) pNIV2870,  $F_{\text{RSV}} \times G_{\text{RSV}}$

15 ELISA were done using 20RG45, a goat anti-RSV serum (Fitzgerald, U.S.A.) to  
coat the microtiter plates and mAb19 anti-F RSV (G.Taylor, Inst. of Animal  
Health, Compton Lab., U.K.) as capture antibody.

20 B) Expression in CHO cells (stable transformants)

All recombinant plasmids were transfected by calcium phosphate coprecipitation  
into CHO-K1 cells, using 20  $\mu\text{g}$  DNA per  $1.25 \times 10^6$  cells. The CHO-K1 cells were  
grown in GMEM-S medium. The GS transfectants were selected by adding 25  $\mu\text{M}$   
25 methionine sulfoximine to the culture medium two days after transfection. After ten  
to fourteen days, resistant colonies were picked and transferred into 96 wells plates.  
Each transformant was then transferred into 24 wells plates and subsequently to 80  
 $\text{cm}^2$  flasks. The GS transformants were assayed for the recombinant products when  
cells reached about 80% confluency. The procedure follows the one described in  
30 Cockett *et al* (Bio/Technology, 1990, 8, 662-667).

ELISA and immunoprecipitation of radiolabelled products were done using the same procedures as the ones described above for the pSFV1 system.

## Results

5

	PSFV1		CHO			
Chimeric products	Expression	Secretion	Expression	Secretion	Size (kDa)	
F <sub>RSV</sub> G <sub>RSV</sub>	+	+	+	+	±130	homochimeric products
F <sub>PIV3</sub> HN <sub>PIV3</sub>	+	-	undetectable			
F <sub>MuV</sub> HN <sub>MuV</sub>	+	-	undetectable			
F <sub>RSV</sub> HN <sub>MuV</sub>	+	+	+	+	± 135	heterochimeric products
F <sub>RSV</sub> HN <sub>PIV3</sub>	+	+	+	+	± 130	
F <sub>PIV3</sub> G <sub>RSV</sub>	+	+	+	+	± 130	
F <sub>PIV3</sub> HN <sub>MuV</sub>	+	+	+	+	± 130	
F <sub>MuV</sub> G <sub>RSV</sub>	+	+	+	+	± 130	
F <sub>MuV</sub> HN <sub>PIV3</sub>	+	+	+	+	± 120	

## Expression in Insect cells

### 10 a) Expression in lepidopteran cells.

The vector pAcUW51 is a shuttle vector for bacteria and lepidopteran cells. A heterologous protein coding sequence can be inserted downstream the baculovirus p10 promoter or either downstream the polyhedrin promoter.

15

The pAcGP67 vector is a shuttle vector for bacteria and lepidopteran cells that contains the gp67 signal sequence upstream a multiple cloning site. A heterologous gene can be inserted in one of the cloning site and will be expressed as a gp67

signal peptide fusion protein under the control of the polyhedrin promoter. The gp67 signal peptide mediates the secretion of the recombinant protein.

5 Either pAcUW51 or pAcGP67 recombinant plasmid can be transfected along with baculovirus linearised DNA into Sf9 cells (Baculogold DNA, PharMingen). This leads to the generation of a recombinant baculovirus stock. The expression of the recombinant heterologous protein is obtained by infecting insect cells with the recombinant baculovirus

10 Plasmid pNIV4132 or plasmid pNIV4136 were transfected with baculovirus linearised DNA into Sf9 cells. Recombinant baculovirus 3546 (derived from cells transfected by pNIV4132) or 5V (derived from cells transfected by pNIV4136) were plaque purified and were used to infect Sf9 or High Five™ cells (Invitrogen). 24h to 72 h post-infection the cells and the spent culture medium have been collected for ELISA and  
15 Western blot analysis.

ELISA were done using anti-RSV goat serum 20RG45 (Fitzgerald) to coat microtiter plates and mAb19 anti-F RSV (G.Taylor) as a capture antibody.

20 Western blots were done using mAb19 anti-F RSV (G.Taylor) or using anti-RSV goat polyclonal serum 20RG45 (Fitzgerald).

The spent medium from cells infected by either baculovirus 3546 or by 5V tested positive in ELISA. The level of expression, depending on the host cell line (SF9 or  
25 High Five), multiplicity of infection, medium (fetal calf serum supplemented or serum free synthetic medium) was at least ten times higher than the one obtained with a recombinant CHO-K1 clone obtained by transfection with pEE14 F<sub>RSV</sub> (1-526) HN<sub>PIV3</sub> (70-572)bis .

30 In addition, the spent medium of the baculovirus infected cells reacted positively in Western blot. A band in the vicinity of 110kDa was present in the immunoblots. These

results confirm the secretion of the chimeric  $F_{RSV}$ - $HN_{PIV3}$  into the medium of Sf9 and High Five cells infected with the recombinant baculoviruses.

#### b) Purification of the recombinant product

5

SF9 cells, adapted to serum free medium, were infected with the plaque purified recombinant baculovirus V5 or 3546. The cells were grown in suspension in 500ml Erlenmeyer flask in SF900II medium (Gibco BRL). The medium from virus infected cells were harvested two days post-infection. The soluble  $F_{RSV}$ - $HN_{PIV3}$  product was purified from the medium of infected cells by immunoaffinity chromatography using an anti-F RSV monoclonal antibody, mAb19. The anti-F monoclonal antibody was coupled to Activated CH Sepharose 4B (Pharmacia) following the manufacturer instructions. The immunoaffinity gel was washed 3 times with 10 bed volumes of buffer A (20mM phosphate buffer pH 6.4, NaCl 150mM) prior to sample loading. After 16 hours at 4°C, the gel was washed with buffer A and the chimeric product was eluted with 100mM phosphoric acid. Eluted protein was neutralized immediately with one tenth of volume of 1M phosphate buffer pH 7.

SDS-PAGE of the immunoaffinity-purified  $F_{RSV}$ - $HN_{PIV3}$  revealed the presence of a major protein band of about 110 kDa. This protein was visualized by Coomassie blue staining of the gel and reacted with the monoclonal antibody anti- $F_{RSV}$  (mAb19) or with the polyclonal serum (20RG45) on immunoblots (Fig25).

#### c) production of polyclonal antibodies

25

In order to obtain specific antibodies, the baculovirus derived  $F_{RSV}$ - $HN_{PIV3}$  protein, purified by immunoaffinity as described above, was used to immunise four BalbC mice and two New Zealand white rabbits. Three sub-cutaneous injections of 20µg/ml/dose/rabbit or 6µg/100µl/dose/mouse were done at three weeks interval. The sera were collected 3 weeks after the second and the third injection and the antibody response was detected using ELISA and Western blots assays.

## 1) ELISA assays

## a) Mice response

The antibody response was followed using a goat anti-RSV serum (2ORG45, Fitzgerald, USA) to coat the microtiter plates and mouse anti-FHN sera as capture antibody. The antigens used were either the  $F_{RSV}$ -Drosophila or CHO derived, the  $F_{RSV}$ -HN<sub>PIV3</sub> expressed in baculovirus and the medium of CHO cells transfected by the pEE14 was used as a negative control.

3 out of 4 mice sera collected after the second injection showed some but low specific response. However, the mice sera collected after the third injection showed a high increase in level of specific antibodies.

## b) Rabbit response

The antibody response was followed using either one of the following ELISA. The antigens were the same as the one used to detect the mice antibody response.

Either a goat anti-RSV serum (2ORG45, Fitzgerald, USA), either a monoclonal antibody directed against the RSV fusion protein (mAb19, Compton Lab, UK) or a monoclonal antibody directed against the PIV3 hemagglutinin-neuraminidase (mAb3285, Behring) were used to coat the microtiter plate and the rabbit anti-HN sera was used as a capture antibody. The first and the second test bleeds generated high specific antibodies.

## 2) Western blot assays

Recombinant  $F_{RSV}$ -CHO-K1 or Drosophila derived,  $F_{RSV}$ -HN<sub>PIV3</sub> baculovirus derived or the CHO-pEE14 spent medium culture were electrophoresed onto a 15% SDS-PAGE and transferred onto a nitrocellulose membrane (Amersham). The rabbit anti-HN sera as well as the mouse anti-HN sera detected specifically either the F protein or the  $F_{RSV}$ -HN<sub>PIV3</sub> chimera.

**Example 2**

- i) Optimization of the codon usage of the nucleic acids sequence coding for the RSV fusion protein lacking the membrane anchor domain linked to the PIV3 hemagglutinin-neuraminidase lacking the signal-anchor domain,  $F_{RSV}$  (1-526) HN<sub>PIV3</sub> (70-572) for the expression in mammalian cells.

A table showing the comparison of the codon usage found in the  $F_{RSV}HN_{PIV3}$  module with the one found in highly expressed human gene can be found in Fig.26. As noted, the most prevalent codons found in the  $F_{RSV}HN_{PIV3}$  module have an A or a T at their third degenerative position, whereas the human prevalent codons have a C or a G. For the improvement of the  $F_{RSV}HN_{PIV3}$  protein expression, the entire coding sequence has been re-engineered to fit at best the human codon usage. The re-engineered sequence was obtained using synthetic long oligonucleotides, polymerase chain reaction (PCR) and conventional cloning procedures.

15

**Re-engineering of the coding sequence of the  $F_{RSV}HN_{PIV3}$  module**

The entire synthetic sequence was recovered by joining three PCR fragments (A, B and C). The general strategy to obtain each PCR fragment is schematically represented in Fig 27. It consists of assembling overlapping long oligonucleotides in a first round amplification. The resulting full size fragment is further amplified using two short primers located on each of its extremities.

20

**Construction of fragment A**

The first PCR fragment, corresponding to 18 bases encoding restriction sites followed by bases 1 to 1269 of the  $F_{RSV}HN_{PIV3}$  followed by 8 bases encoding restriction sites, was obtained by PCR assembly of 18 overlapping oligonucleotides (Fig 28). This fragment has been inserted in the pCRIITOPPO cloning vector (Invitrogen). After sequencing the fragment, it was retrieved from the pCRIITOPPO vector by a *XbaI* and *BsrGI* digestion and inserted into the corresponding sites of pNIV4120. The module corresponding to  $F_{RSV}HN_{PIV3}$  with bases 1 to 1264 humanized was then retrieved by an *XbaI* and *EcoRI* digestion and inserted into the corresponding sites of pEE14 (Fig.29) generating pEE14x $F_{RSV}humHN_{PIV3}$ .

30

### Construction of fragment B

The second PCR fragment B corresponding to 13 bases encoding unique restriction sites followed by bases 1264 to 2136 of  $F_{RSV}HN_{PIV3}$  was obtained by assembling 10 oligonucleotides whose sequences can be found in Fig.30. This fragment has been inserted in the pCRIITOPPO vector and sequenced. This fragment has been recovered by a *BsrGI* and *KpnI* digestion.

### Construction of fragment C

The third PCR fragment corresponding to bases 2023 to 3090 followed by 6 extra bases encoding an *EcoRI* site has been assembled starting from the 15 oligonucleotides shown in Fig 31. This fragment has been inserted in the pCRIITOPPO cloning vector and sequenced. This fragment has been retrieved by a *KpnI* and *EcoRI* digestion (Fig 31).

### Construction of the entire coding sequence

The entire  $F_{RSV}HN_{PIV3}$  codon optimized coding sequence has been obtained by assembling fragment A, B, C as shown in Fig.32. pNIV4120 in which the PCR fragment A has replaced the original sequence (see Fig.29) was digested by *BsrGI* and *EcoRI*. The original sequence was eliminated and replaced by the *BsrGI*-*KpnI* fragment B and the *KpnI*-*EcoRI* fragment C. The codon optimized module was retrieved from the PCRITOPPO vector by a *XbaI* and an *EcoRI* and inserted in the corresponding sites of the pEE14 vector. The resulting plasmid, pEE14 $F_{RSV}$  hum $HN_{PIV3}$  hum, encodes for the entire humanized coding sequence. The humanized  $F_{RSV}HN_{PIV3}$  nucleic acids sequence is shown in Fig. 33.

### Expression in CHO-K1 cells

The recombinant pEE14  $F_{RSV}$  hum $HN_{PIV3}$  (see construction of fragment A, above, or recombinant pEE14 $F_{RSV}$  hum $HN_{PIV3}$  hum see construction of the entire coding sequence, above) was transfected using the FuGene reagent (Boeringer Mannheim), using 5  $\mu$ g DNA per  $1.25 \times 10^6$  cells. The CHO-K1 cells were grown in GMEM-S medium. The GS transfectants were selected by adding 25  $\mu$ M methionine sulfoximine to the culture medium two days after transfection. After ten to fourteen

days, resistant colonies were picked and transferred into 96 wells plates. Each transformant was then transferred into 24 wells plates and subsequently to 80 cm<sup>2</sup> flasks. The GS transformants were assayed for the recombinant product when cells reached about 80% confluency. The procedure follows the one described in Cockett  
5 *et al* (Bio/Technology, 1990, 8, 662-667). Alternatively, the expression was evaluated three to five days after the addition of sodium butyrate (2mM) in the cell culture.

To compare the expression level to that of the non humanized F<sub>RSV</sub>HN<sub>PIV3</sub>, ELISA  
10 assays were done, using 20RG45, a goat anti-RSV serum (Fitzgerald, U.S.A.) to coat the microtiter plates and mAb19 anti-F RSV (G. Taylor, Inst. of Animal Health, Compton Lab, U.K.) as capture antibody. The expression level was estimated using a purified Fa<sub>RSV</sub> expressed in the Drosophila system.

15 The level of expression of the non-humanized expressed product by pEE14F<sub>RSV</sub>HN<sub>PIV3</sub> didn't exceed 0.03 mg/L and 0.1 mg/L when sodium butyrate was added to the culture medium. The level of expression of the partially humanized product expressed by pEE14F<sub>RSV</sub> humHN<sub>PIV3</sub>, reached 1 mg/L and up to 3 mg/L when sodium butyrate was added in the culture medium. The humanization  
20 of the sequence coding for amino acids 1-423 of the 1029 amino acids thus enhanced the level of expression up to 30 fold (see Figure 34a).

The level of expression of the entirely humanized product expressed by pEE14F<sub>RSV</sub> humHN<sub>PIV3</sub>hum was at least of 2 mg/L and reached up to 50 mg/L when sodium  
25 butyrate was added in the culture medium. The humanization of the entire coding region of F<sub>RSV</sub>HN<sub>PIV3</sub> thus enhanced the level of expression of at least 200 to 500 fold (see Figure 34b).

ii) Optimization of the codon usage of the nucleic acids sequence coding for the  
30 mumps virus (MuV) fusion protein lacking the membrane anchor domain linked to the measles virus (MV) lacking the signal-anchor domain, F<sub>MuV</sub> (1-482) H<sub>MV</sub> (59-617) for the expression in mammalian cells.

A table showing the comparison of the codon usage found in the  $F_{MuV}H_{MV}$  module with the one found in highly expressed human gene can be found in Fig.35. As it can be seen, the codon usage frequencies of this chimerical gene is quite different from those prevalent in the human genome. For the improvement of the  $F_{MuV}H_{MV}$  protein expression, the entire coding sequence has been re-engineered to fit at best the human codon usage. The re-engineered sequence was obtained using synthetic long oligonucleotides, polymerase chain reaction (PCR) and conventional cloning procedures.

#### Re-engineering of the coding sequence of the $F_{MuV}H_{MV}$ module

The entire synthetic sequence was recovered by joining four PCR fragments (A, B, C and D). The general strategy to obtain each PCR fragment is schematically represented in Fig 36. It consists of assembling overlapping long oligonucleotides in a first round amplification. The resulting full size fragment is further amplified using two short primers located on each of its extremities.

#### Construction of fragment A

The first PCR fragment, corresponding to 13 bases specifying restriction sites and a Kozak consensus motif followed by bases 1 to 1026 of the  $F_{MuV}H_{MV}$  was obtained by PCR assembly of 12 overlapping oligonucleotides (Fig 37). This fragment has been inserted in the pCRIITOPPO cloning vector (Invitrogen). After sequencing the fragment, it was retrieved from the pCRIITOPPO vector by a *XbaI* and *TspRI* digestion and a 963 bp fragment was further purified, leading to fragment A.

#### Construction of fragment B

The second PCR fragment B corresponding to bases 965 to 1712 of  $F_{MuV}H_{MV}$  was obtained by assembling 9 oligonucleotides whose sequences can be found in Fig.38. After its insertion into the pCRIITOPPO vector and its sequencing, this 785 bp fragment has been recovered by a *TspRI* and *AvaI* digestion.

### Construction of fragment C

The third PCR fragment C corresponding to bases 1712 to 2485 has been assembled starting from the 11 oligonucleotides shown in Fig 39. It has been inserted in the pCRIITOPPO cloning vector and sequenced. This 774 bp fragment has been retrieved  
5 by an *AvaI* and *Apal* digestion.

### Construction of fragment D

The fourth PCR fragment D corresponding to bases 2485 to 3139 followed by 8 bp specifying a unique restriction site has been assembled starting from the 8  
10 oligonucleotides shown in Fig 40. This fragment has been inserted in the pCRIITOPPO vector and sequenced. A 657 bp fragment has been recovered after an *Apal* and *EcoRI* digestion.

### Construction of the entire coding sequence

15 The entire  $F_{MuV}H_{MV}$  codon optimised coding sequence has been obtained by assembling fragment A, B, C, D and inserting the module digested by *XbaI* and *EcoRI* into the corresponding sites of the pEE14 vector (Fig. 41). The resulting plasmid, pEE14 $F_{MuV}humH_{MV}hum$ , encodes for a humanised sequence coding for aa 1-482 of the mumps virus fusion protein followed by aa 59-617 of the measles  
20 virus. The humanised and original  $F_{MuV}H_{MV}$  nucleic and amino acids sequences are shown in Fig. 42.

### iii) Purification and analysis of FHN expressed in CHO-K1

#### 25 a) Purification

CHO cell line expressing secreted recombinant FHN was cultivated in cell factories in G-MEM medium supplemented with 2% FCS, in presence or absence of 1% Butyrate Na. FHN was purified by immunoaffinity chromatography by loading spent culture  
30 medium onto a Mab19-sepharose column as described using the same experimental conditions.

When expressed in absence of Butyrate Na, purified FHN migrated on SDS-PAGE, in heating and reducing conditions, mainly as a band of 110 kDa. In contrast, FHN is visualized as a triplet of 110, 120 and 130 kDa when CHO cells are cultivated with butyrate. Heating has a more drastic effect than reducer on the FHN electrophoretic migration. Indeed, high molecular weight species are clearly detected in the preparation when electrophoresis proceeded without heating suggesting the presence of FHN aggregates or oligomers. These aggregates did not seem to be contaminated by CHO proteins. Antibodies directed to CHO proteins did not specifically recognize on Western blot any bands. Glycan analysis was performed using several lectins specific for different carbohydrate moieties. Surprisingly, FHN did not carry sialic acids or high-mannose structures but carbohydrates of galactose-acetyl-galactosamine type characteristic of hybrid N- and/or O-glycosylations.

N-terminal microsequence analysis showed mainly the presence of F1 subunit in bands of 110-130kDa. The F2 N-terminal amino acid sequence detected in bands of lower and higher molecular weight indicated that some purified FHN molecules are present under a F0 form (non mature F).

The presence of aggregates or oligomers in the FHN preparations was confirmed by gel filtration analysis and proteins were detected by laser-light scattering. Whatever the culture conditions (butyrate or not), between 50 and 65% of FHN populations displayed a molecular weight higher than  $10^6$  Da demonstrating that FHN is aggregated. 5 to 15% has a molecular weight ranging from 400 to 900 kDa whereas 30 to 35% is monomeric FHN.

#### b) Serum immunoglobulin analysis.

##### Immunisation protocol

The  $F_{RSV}HN_{PIV3}$  protein was purified from the spent medium culture of the CHO-K1 cells transfected by the recombinant pEE14  $F_{RSV}humHN_{PIV3}hum$  by immunoaffinity chromatography as described (Purification of the recombinant product expressed in baculovirus recombinant infected SF9 cells). The product was injected in 7 groups of Balb C1 mice as described in the following table 1.

**Humoral response directed against the FHN protein**

The humoral response directed against the FHN protein was determined. To this end, ELISA plates were coated with immunoaffinity purified FHN protein.

5

**Total IgG (Fig 43)**

To detect specific anti-FHN total IgG, ELISA plates were coated with 200ng of immunoaffinity purified FHN protein, plates were then saturated and dilutions of the mice second bleed sera were then applied. Total IgG were detected using a biotinylated serum directed against mouse IgG.

10

**IgG1 (Fig 44)**

To detect specific anti-FHN IgG1, ELISA plates were coated with 100ng of immunoaffinity purified FHN protein, plates were then saturated and dilutions of the mice second bleed sera were then applied. IgG1 were detected using a biotinylated serum directed against mouse IgG1.

15

**IgG2a (Fig 45)**

To detect specific anti-FHN IgG2a, ELISA plates were coated with 100ng of immunoaffinity purified FHN protein, plates were then saturated and dilutions of the mice second bleed sera were then applied. IgG2a were detected using a biotinylated serum directed against mouse IgG2a.

20

The titer of each sera was determined and a mean titer for each group was calculated and is reported in table 2. These experiments show that the FHN antigen by itself or formulated with adjuvant (group 1 to 3), stimulates a specific humoral response. Indeed, no anti-FHN antibodies are generated in the untreated mice group (group 5) or in the group immunised solely with the adjuvant (group 4). The group 1 (and group 4) adjuvant was 3D-MPL and QS21 formulated with cholesterol containing liposomes as described in WO 96/33739; the group 2 adjuvant was alum.

30

The IgG1/IgG2a ratio indicates the Th1 or Th2 orientation of the immune response; (Table2), a protective response against both the RSV or the PiV3 should tend toward

the Th1 type, that is a low IgG1/IgG2a ratio. In this regard, the responses generated with the FHN formulated in the presence of the 3D-MPL + QS21 adjuvant appears to be the more promising one.

# 5 Table 1: Experimental procedures

Immunogenicity FHN in mice

Group	n	Vol ( $\mu$ l)	route	Antigen		Immuno- stimulants	buffer	preservative
				nature	dose ( $\mu$ g)			
1	12	2x50	IM	FHN	2	3D-MPL/ QS21	PBS mod pH 7.4	thiomersal low (1 $\mu$ g/ml)
2	12	2x50	IM	FHN	2	Al(OH) <sub>3</sub>	PBS mod pH 7.4	thiomersal low (1 $\mu$ g/ml)
3	12	2x50	IM	FHN	2	/	PBS mod pH 7.4	thiomersal low (1 $\mu$ g/ml)
4	12	2x50	IM	/	/	3D-MPL/ QS21	PBS mod pH 7.4	thiomersal low (1 $\mu$ g/ml)
5	12	/	/	untreated	/	/	/	/
6	12	2x30	INA	RSV live		/	/	/
7	12	2x30	INA	PIV-3 live		/	/	/

IM=intra-muscular

INA=intra-nasal

Antigen	cc. $\mu$ g/ml	Buffer
RSV live	6.2 logPFU/ml	
PIV-3 live	6.7 logPFU/ml	
FHN	120 (2.5ml)	PBS pH 7.3

## Time schedule:

- 5 Injection 1 = Day 0  
Injection 2 = Day 28  
First Bleed = Day 28  
Second bleed = Day 42

Antigen	cc. µg/ml	Buffer
RSV live	6.2 logPFU/ml	
PIV-3 live	6.7 logPFU/ml	
FHN	120 (2.5ml)	PBS pH 7.3

## Time schedule:

- 5 Injection 1 = Day 0  
Injection 2 = Day 28  
First Bleed = Day 28  
Second bleed = Day 42

**Table 2: Serum antibody response against FHN.**

The total IgG, IgG1 and IgG2a was determined for each mouse sera. A mean titer for each group was then calculated and is reported in the table.

group n°	Immunogen	Total IgG	IgG1	IgG2a	IgG1/IgG2a
1	FHN + 3D-MPL/QS21	1182000	109800	305500	0.36
2	FHN + Alum	182200	127100	4429	28.7
3	FHN	44990	22760	1941	11.73
4	adjuvant=from group 1	49	32	ND	ND
5	untreated	52	ND	ND	ND
6	Live RSV	12840	748	2718	0.27
7	Live PiV3	10860	2758	2320	1.19

ND=undetermined, the titer being too low

5

### References

- Haas J., Park E-C. and Seed B., Codon usage limitation in the expression of HiV-1 envelope glycoprotein, *Current Biology*, 1996, 6, n°3, 315-325.
- 10 Kim C. H., Oh Y. and Lee T.H., Codon optimization for high-level expression of human erythropoietin (EPO) in mammalian cells, *Gene*, 1997, 199, 293-301.
- Zolotukhin S., Potter M. Hauswirth W.W. Guy J. and Muzyczka N. A Humanized green fluorescent protein cDNA adapted for high level expression in mammalian cells. *J. of Virology*, July 1996, 70, n°7, 4646-4654.

**Claims**

1. A process for preparing a heterochimeric protein or an immunogenic derivative thereof comprising an immunogenic fragment of the fusion (F) protein of RSV, PIV1, PIV2, PIV3, MV or MuV and an immunogenic fragment of the attachment (G, HN or H) protein of RSV, PIV1, PIV2, PIV3, MV or MuV which process comprises expressing recombinant DNA encoding the heterochimeric protein or immunogenic derivative thereof in CHO cells and recovering the protein.
2. A process according to claim 1 wherein at least one non-preferred or less preferred codon in a natural gene or DNA encoding the said heterochimeric protein or immunogenic fragment thereof has been replaced by a preferred codon encoding the same amino acid.
3. A heterochimeric protein or an immunogenic derivative thereof comprising an immunogenic fragment of the fusion (F) protein of RSV, PIV1, PIV2, PIV3, MV or MuV and an immunogenic fragment of the attachment (G, HN or H) protein of RSV, PIV1, PIV2, PIV3, MV or MuV, with the proviso that where one of the immunogenic fragments is derived from RSV F, RSV G or PIV3 F, PIV3 HN, the other of the immunogenic fragments is derived from MuV F, MuV HN, MV F, MV H, PIV1 F, PIV1 HN, PIV2 F or PIV2 HN.
4. A process for preparing a heterochimeric protein or immunogenic derivative thereof as claimed in claim 3 which process comprises expressing recombinant DNA encoding the heterochimeric protein or immunogenic derivative thereof in either one of; CHO cells or insect cells and recovering the protein.
5. A protein according to claim 3 wherein the immunogenic fragment of the F protein is lacking the membrane anchor domain at its C-terminal end.
6. A protein according to claims 3 or 5 wherein the immunogenic fragment of the G, HN or H protein is lacking the signal/anchor domain at its N-terminal end.

7. A protein according to any one of claims 3, 5 or 6 which is linked *via* an amino acid in the C-terminal part of the immunogenic fragment of the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV to an amino acid in the N-terminal part of the immunogenic fragment of the G protein of RSV or the HN protein of PIV1, PIV2, PIV3, MuV or the H protein of MV.
8. A protein according to any one of claims 3, 5, 6 or 7 which commences at its N-terminal end with a signal sequence from the F protein of RSV, PIV1, PIV2, PIV3, MV or MuV.
9. A protein according to any one of claims 3, 5, 6 or 7 which commences at its N-terminal end with a signal sequence from TPA.
10. A protein according to any one of claims 3 or 5 to 8 which comprises a ubiquitin leader sequence.
11. A protein according to any one of claims 3 or 5 to 9 which comprises a polyhistidine tail.
12. A protein according to claim 10 or 11 which comprises a cleavage site for cleaving off the ubiquitin leader sequence and/or the polyhistidine tail.
13. A heterochimeric protein according to any one of claims 3 or 5 to 11 which is selected from the group consisting of:
- Fs<sup>+</sup>a<sup>-</sup>RSVxHNs<sup>-</sup>a<sup>-</sup>MuV;  
Fs<sup>+</sup>a<sup>-</sup>PIV3 x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup>MuV;  
Fs<sup>+</sup>a<sup>-</sup>MuV x Gs<sup>-</sup>a<sup>-</sup>RSV; or  
Fs<sup>+</sup>a<sup>-</sup>MuV x HN<sup>-</sup>s<sup>-</sup>a<sup>-</sup>PIV3, or  
an immunogenic derivative thereof.
14. A heterochimeric protein according to any one of claims 3 or 5 to 11 which is selected from the group consisting of:

Fs<sup>+</sup>a<sup>-</sup> MuV x Hs<sup>+</sup>a<sup>-</sup> MV; or  
Fs<sup>+</sup>a<sup>-</sup> RSVx HNs<sup>+</sup>a<sup>-</sup> PIV1, or  
Fs<sup>+</sup>a<sup>-</sup> RSVx HNs<sup>+</sup>a<sup>-</sup> PIV2, or  
an immunogenic derivative thereof.

5

15. A heterochimeric protein which is:

Fs<sup>+</sup>a<sup>-</sup> (1-526) RSV x HNs<sup>+</sup>a<sup>-</sup> (70-572) PIV3,  
Fs<sup>+</sup>a<sup>-</sup> (1-492) PIV3 x Gs<sup>+</sup>a<sup>-</sup> (69-298) RSV,  
Fs<sup>+</sup>a<sup>-</sup> (1-526) RSV x HNs<sup>+</sup>a<sup>-</sup> (70-572) PIV3 bis,  
10 Fs<sup>+</sup>a<sup>-</sup> (1-526) RSV x HNs<sup>+</sup>a<sup>-</sup> (70-572) PIV3 ent his, or  
sTPA (1-21) UB (1-74) ent Fs<sup>+</sup>a<sup>-</sup> (24-526) x HN s<sup>+</sup>a<sup>-</sup> (70-572) PIV3, or  
an immunogenic derivative thereof.

15

16. Recombinant DNA encoding a heterochimeric protein or an immunogenic  
derivative thereof according to any one of claims 3 or 5 to 15.

17. Recombinant DNA according to claim 16 in which at least one non-preferred  
or less preferred codon in the DNA has been replaced by a preferred codon  
encoding the same amino acid.

20

18. DNA which hybridises under conditions of high stringency with the DNA of  
claim 16 or 17.

19. An expression vector comprising recombinant DNA according to claims 16 to  
18.

20. A host transformed with DNA according to any one of claims 16 to 18 or with  
a vector according to claim 19.

25

21. A host according to claim 20 which is a CHO cell.


22. A host according to claim 21 which is an insect cell.

23. A vaccine composition comprising a protein according to any one of claims 3 or 5 to 13 or an immunogenic derivative thereof in admixture with a pharmaceutically acceptable carrier.
24. A vaccine composition according to claim 23 further comprising 3D
- 5 Monophosphoryl lipid A and/or QS-21.
25. A vaccine composition according to claims 23 or 24 wherein the carrier is an oil-in-water emulsion.
26. A heterochimeric protein or an immunogenic derivative thereof according to any one of claims 3 or 5 to 15 for use in medicine.
- 10 27. A process for the production of a heterochimeric protein according to any one of claims 3 or 5 to 15 which process comprises expressing recombinant DNA encoding said protein or immunogenic fragment thereof in a host cell and recovering the protein.
28. A method of treating a human or animal susceptible to paramyxoviridae viral
- 15 infections comprising administering an effective amount of a vaccine according to any one of claims 23 to 25.
29. Use of a protein or an immunogenic derivative thereof according to any one of claims 3 or 5 to 15 in the manufacture of a medicament for use in the treatment of respiratory disorders.

1/73

Fig. 1


pNIV2819



2/73

Fig. 2


pNIV2820



3/73

Fig. 3

pNIV2841



4/73

Fig. 4

## A) Synthetic adaptators


5' C ATG AAT GAT CAA GGC TTG AGC AA 3'

TTA CTA GTT CCG AAC TCG TTA GTC [SEQ ID NO: 1]


BspHI

BbsI

## B) pNIV4102



## C) pNIV4104



5/73

Fig. 5

## A) Synthetic adaptators

5' C ATG AAC AAT GAG TTT ATG GAA GTT ACA GAA AAG ATC CAA  
 TTG TTA CTC AAA TAC CTT CAA TGT CTT TTC TAG GTT

BspHI


ATG GCA TCG GAT ATT AT 3'

TAC CGT AGC CTA TAA TATA


[SEQ ID NO: 2]

AseI

## B) pNIV4109



## C) pNIV4110



6/73

Fig. 6

## A) Synthetic adaptators


5' GAT CTA GAA GAG TCA AAA GAA TGG ATA AGA AGG TCA AAT CAA  
 AT CTT CTC AGT TTT CTT ACC TAT TCT TCC AGT TTA GTT

BglII


AAA CTA GAT TCC ATT GGA AAT TGG CAT CAA TCT AGC ACC 3'  
 TTT GAT CTA TGG TAA CCT TTA ACC GTA GTT AGT TCG TGG CAGT G  
 MaeIII

[SEQ ID NO: 3]

## B) pNIV4103



## C) pNIV4106



7/73

Fig. 7


## A) Synthetic adaptators

5'G ATC TAG AAG AGT CAA AAG AAT GGA TAA GAA GGT CAA ATC  
 ATC TTC TCA GTT TTC TTA CCT ATT CTT CCA GTT TAG  
 BglII


AAA AAC TAG ATT CCA TTG GAA ATT GGC ATC AAT CTA GCA CCA  
 TTT TTG ATC TAA GGT AAC CTT TAA CCG TAG TTA GAT CGT GGT

CAA ATG ATC AAG GCT TGA GCA A 3'  
 GTT TAC TAG TTC CGA ACT CGT TAGTC [SEQ ID NO: 4]  
 BbsI

## B) pNIV4117



## C) pNIV4118



8/73


Fig. 8

## A) Synthetic adaptators


5' GAA TGC CGT TAA ATA CAT CAA GAG AGT AAC CAT CAA  
 A CGT CTT ACG GCA ATT TAT GTA GTT CTC TCA TTG GTA GTT  
 PstI

CTC CAT CGG TCT CAG TAA GTT CTA AA 3'  
 GAG GTA GCC AGA GTC ATT CAA GAT TTC AGT [SEQ ID NO: 5  
 MaeIII

## B) pNIV4113



## C) pNIV4114




9/73

Fig. 9


## A) Synthetic adaptators

5' GTAAGTTCTAAA 3'  
CAAGATTTTAA [SEQ ID NO: 6]  
BsaI EcoRI

## B) pNIV4115





## C) pNIV4116



10/73

Fig. 10


## A) pNIV4102

C) pEE14 F<sub>s</sub><sup>+</sup>a<sup>-</sup> RSV x HN s<sup>-</sup>a<sup>-</sup> MuV



11/73

Fig. 11

## A) pNIV4105





## B) pNIV4109

C) pEE14  $Fs^+a^-$  RSV x  $HN s^-a^-$  PIV3

12/73

Fig. 12



## A) pNIV4103

B) pEE14 F<sup>s</sup>+a<sup>-</sup> PIV3 x G<sup>s</sup>-a<sup>-</sup> RSV

13/73

Fig. 13



## A) pNIV4117

B) pEE14 F<sub>s</sub><sup>+</sup>a<sup>-</sup> PIV3 x HN s<sup>-</sup>a<sup>-</sup> MuV

14/73

Fig. 14



## A) pNIV4113

B) pEE14  $Fs^+a^-MuV \times Gs^+a^-RSV$ 

15/73

Fig. 15


## A) pNIV4115

B) pEE14  $Fs^+a^-MuV \times HNs^-a^-PIV3$ 


16/73

Fig. 16

## A) pNIV2857





## B) pNIV2870



17/73

Fig. 17

## A) pNIV2857

B) pEE14  $Fs^+a^-$  RSV x  $Gs^+a^-$  RSV

18/73

Fig. 18

## A) Synthetic adaptators

5' C ATG AAC AAT GAG TTT ATG GAA GTT ACA GAA AAG ATC CAA  
 TTG TTA CTC AAA TAC CTT CAA TGT CTT TTC TAG GTT



BspHI

ATG GCA TCG GAT ATT AT 3'  
 TAC CGT AGC CTA TAA TAT A

[SEQ ID NO: 7]

AseI

## B) pNIV4120

C) pEE14 F s<sup>+</sup> a<sup>-</sup> RSV xHN s<sup>-</sup> a<sup>-</sup> PiV3bis

19/73

Fig. 19


## A) Synthetic adaptators

PstI 5'GT GAC GAT GAC GAT AAG CAT CAT CAT CAT CAT CAT TAG  
 ACGTC ACA CTG CTA CTG CTA TTC GTA GTA GTA GTA GTA GTA ATC



GGATCCGCATG 3'

CCTAGGC SphI [SEQ ID NO: 8]

## B) pNIV3340




## C) pNIV4120

C) pEE14 F s<sup>+</sup> a<sup>-</sup> RSV xHN s<sup>-</sup> a<sup>-</sup> PiV3 enthis

20/73

Fig. 20

## A) pNIV3475




## B) Synthetic adaptators

5' CT AGC ATG CAG ATC TTC GTC AAG ACG TTA ACC GGT AAA ACC  
 NheI G TAC GTC TAG AAG CAG TTC TGC AAT TGG CCA TTT TGG

ATA ACC 3' XbaI  
 TAT TGG ATCT

[SEQ ID NO: 9]


## C) pNIV4122



21/73

Fig. 21

## A) pNIV4122



## B) Synthetic adaptators


5' TTA AGA CTA AGA GAC GAT GAC GAT AAG TCC AGT CAA AAC  
 AflII CT GAT TCT CTG CTA CTG CTA TTC AGG TCA GTT TTG

ATC ACT GAA GAA TTT TAT CAA TCA ACA TGC AGT GCA GTC AGC  
 TAG TGA CTT CTT AAA ATA GTT AGT TGT ACG TCA CGT CAG TCG

AAA GGC TAT CTT AGT GCT CTA AGA ACT GGT TGG TAT A 3' SpeI  
 TTT CCG ATA GTT TCT CGA GAT TCT TGA CCA ACC ATA TGA TC

[SEQ ID NO: 10]


## C) pNIV4123




22/73

Fig. 22


## A) pNIV4123




## B) pNIV4120



## C) pNIV4124




## D) pEE14 sTPA UBI EN Fs'a' RSV x HN s'a' Piv3




23/73

Fig. 23

## A) pNIV4120




## B) pNIV4132



24/73

Fig. 24

## A) pNIV4120




## B) Synthetic adaptators

5' GAT CAA AAC ATC ACT GAA GAA TTT TAT CAA TCA ACA TGC  
 BamHI TT TTG TAG TGA CTT CTT AAA ATA GTT AGT TGT ACG

AGT GCA GTC AGC AAA GGC TAT CTT AGT GCT CTA AGA ACT  
 TCA CGT CAG TCG TTT CCG ATA GAA TCA CGA GAT TCT TGA

GGT TGG TAT A 3' SpeI  
 CCA ACC ATA TGA TC [SEQ ID NO: 11]

## C) pNIV4136




25/73

**Fig 25: SDS-PAGE (reduced conditions) of the F<sub>RSV</sub>HN<sub>PIV3</sub> protein purified by immunoaffinity from the spent culture medium of the recombinant baculovirus 3546.**

kDa: molecular weight marker

A: Coomassie blue staining

B: Western blot revealed by a goat polyclonal anti-RSV serum 20 RG45



SUBSTITUTE SHEET (RULE 26)


26/73

Fig. 26: Codon usage of F<sub>RSV</sub>HN<sub>PIV3</sub> and highly expressed human genes (hum high exp) showing frequencies (x100) of the individual codons for each of the degenerately encoded amino acids, and the most prevalent codon in bold.

Ala	GCG	17	3	Gln	CAG	<b>88</b>	32	Leu	TTG	6	10	Ser	AGT	10	21
	A	13	<b>51</b>		C	12	<b>68</b>		A	2	<b>39</b>		C	34	16
	T	17	35						CUG	<b>58</b>	6		TCG	9	7
	C	<b>53</b>	11						A	3	23		A	5	<b>27</b>
Arg	AGG	18	28	Glu	GAG	75	22		T	5	11		T	13	17
	A	10	<b>56</b>		A	25	<b>77</b>		C	26	12		C	28	13
	CGG	21	0	Gly				Lys	AAG	<b>82</b>	34	Thr	ACG	15	6
	A	6	10		GGG	24	22		A	18	<b>66</b>		A	14	<b>56</b>
	T	7	5		A	14	<b>37</b>						T	14	26
	C	<b>37</b>	0		T	12	22	Phe	TTT	20	<b>63</b>		TAT	26	<b>79</b>
Asn	AAT	22	<b>62</b>	His	CAT	21	<b>86</b>		C	<b>80</b>	37		C	74	21
	C	<b>78</b>	38		C	<b>79</b>	14	Pro	CCG	17	12	Val	GTG	<b>64</b>	19
Asp	GAT	25	<b>66</b>	Ile	ATA	5	<b>49</b>		A	16	<b>51</b>		A	5	<b>41</b>
	C	<b>75</b>	34		T	18	30		T	19	26		T	7	23
Cys	TGT	32	<b>62</b>		C	<b>77</b>	20		C	<b>48</b>	12		C	25	17
	C	<b>68</b>	38												

27/73

**Fig. 27: Schematic diagram of the PCR synthesis of each fragment showing unique restriction sites along the sequence (black dots) and restriction sites (A and B) that allow retrieval of the full size fragment from the cloning vector.**



28/73

**Fig. 28: Sequence of the 18 oligonucleotides from which PCR fragment A was generated.**




- 1) olfhum1.seq, bases 1-90 of  $F_{RsvHN_{Piv3}}$ , homologous to mRNA  
5' cccTCTAGAGGATCCACCATGGAGCTGCTGATtttaAAGACCAACGCCATCACCGCCATCCTG  
GCCGCGGTGACCCTCTGCTTCGCGTCC
- 2) olfhum2.seq, bases 75-165 of  $F_{RsvHN_{Piv3}}$ , inverse complementary to mRNA  
5' CCTCAGCGCGCTCAGGTAGCCCTTGCTGACAGCagaGCAGGTGGACTGGTAGAACTCCTCGGTG  
ATGTTCTGGCTGGACGCGAAGCAGAGG
- 3) olfhum3.seq, bases 150-240 of  $F_{RsvHN_{Piv3}}$ , homologous to mRNA  
5' CCTGAGCGCGCTGAGGACGGGGTGGTACACTAGtGTGATCACCATCGAGCTGAGCAACATCAAG  
GAGAACAAGTGCAACGGCACCGACGCC
- 4) olfhum4.seq, bases 225-310 of  $F_{RsvHN_{Piv3}}$ , inverse complementary to mRNA  
5' GCATCAGCAGCTGCAGCTCGGTCACGGCGCTCTTGTACTTGTCCAGCTCCTGCTTGATCAGCTT  
CACCTTGGCGTCGGTGCCGTTG
- 5) olfhum5.seq, bases 295-397 of  $F_{RsvHN_{Piv3}}$ , homologous to mRNA  
5' CTGCAGCTGCTGATGCAGAGCACCCCCGCCACCAACAACagaGCCAGGCGCGAGCTGCCCAGGT  
TCATGAACTACACCCTCAACAACACCAAGAACACCAACG
- 6) olfhum6.seq, bases 378-496 of  $F_{RsvHN_{Piv3}}$ , inverse complementary to mRNA  
GGTGCAGGACCTTGGACACCGCGATGCCGCTGGCGATGGCGGAGCCCACGCCCAGCAGGAAGCCCA  
GGAAGcGcctCTTgCgCTTCTTGCTCAGGGTCACGTTGGTGTCTTGGTGTG
- 7) olfhum7.seq, bases 480-561 of  $F_{RsvHN_{Piv3}}$ , homologous to mRNA  
5' GTCCAAGGTCCTGCACCTGGAGGGGGAGGTGAACAAGATCAAGAGCGCCCTGCTCTCCACCAAC  
AAGGCGGTGGTCAGCCTG
- 8) olfhum8.seq, bases 543-633 of  $F_{RsvHN_{Piv3}}$ , inverse complementary to mRNA  
5' GGGGAGCAatTGCTTGTCGATGTAGTTCTTGAGGTCCAGCACCTTGCTGGTCAGCACGCTCACG  
CCGTTGGACAGGCTGACCACCGCCTTG
- 9) olfhum9.seq, bases 609-676 of  $F_{RsvHN_{Piv3}}$ , homologous to mRNA  
5' CTACATCGACAAGCAatTGCTCCCCATCGTGAACAAGCAGtcCTGCAGCATCTTAACATTGAG  
ACCG
- 10) olfhum10.seq, bases 653-732 of  $F_{RsvHN_{Piv3}}$ , inverse complementary to mRNA  
5' GCTGAACTCCCTGGTGATCTCCAGCAGCCTGTTGTTCTTCTGCTGGAACCTCGATCACGGTCTCA  
ATGTTAGAGATGCTGC

29/73

- 11) olfhum11.seq, bases 714-787 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5' GATCACCAGGGAGTTCAGCGTGAACGCgGGcGTcACCACCCCGGTGAGCACCTACATGCTGACC  
AACAGCGAGC
- 12) olfhum12.seq, bases 768-846 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse complementary  
to mARN  
5' GTTGGACATaAGCTTCTTCTGGTCGTTGGTGATGGGCATGTCGTTGATCAGGGACAGCAGCTCG  
CTGTTGGTCAGCATG
- 13) olfhum13.seq, bases 825-916 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5' CCAGAAGAAGCTtATGTCCAACAACGTGCAGATCGTGCGCCAGCAGAGCTACagCATCATGagC  
ATCATCAAGGAGGAGGTGCTGGCCTACG
- 14) olfhum14.seq, bases 900-990 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse complementary  
to mARN  
5' GGTGGTGCACAGGGGGGAGGTGTGCAGCTTCCAGCAGGGGGTGTGTCGATCACGCCGTACAGGGGC  
AGCTGCACCACGTAGGCCAGCACCTCC
- 15) olfhum15.seq, bases 975-1065 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5' CCCCCTGTGCACCACCAACACCAAGGAGGGCTCCAACATCTGCCTGACCCGCACCGACCGGGGC  
TGGTACTGCGACAACGCCGGCTCCGTG
- 16) olfhum16.seq, bases 1048-1133 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mARN  
5' CTGTTTCATGGTGTGTCGAGAACACGCGGTTGGACTGCACCTTGCAGGTCTCCGCCAGGGGGGAAGA  
AGGACACGGAGCCGGCGTTGTC
- 17) olfhum17.seq, bases 1116-1210 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5' CTGCGACACCATGAACAGCCTGACCCTGCCAGCGAGGTGAACCTCTGCAACATCGACATCTTC  
AACCCCAAGTACGACTGCAAGATtATGacctcc
- 18) olfhum18.seq, bases 1195-1295 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mARN  
gggaattctgtacacttggtcttgccgtagcaggacacgatggcgcccagggaggtgatcacggag  
ctgctcacgtcgggtcttgagggtCATAATCTTGCAG

[SEQUENCES ABOVE ARE SEQ ID NOS: 12 to 29, respectively]

30/73

**Fig. 29: Construction of pEE14 F<sub>RSV</sub>humHN<sub>PIV3</sub>.****a) PCR fragment A****b) pNIV4120 +PCR fragment A****c) pEE14 F<sub>RSV</sub>hum HN<sub>PIV3</sub>**

31/73

**Fig. 30: Sequence of the 10 oligonucleotides from which PCR fragment B was generated.**

- 1) olfhnnum19.seq, bases 1269-1353 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'cggcaagaccaagtgtacagcctccaacaagaaccgcgcatcatcaagaccttctccaacgg  
ctgcgactacgtgtccaacaag 3'
- 2)olfhnnum20.seq, bases 1336-1428 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary of mARN  
5'cttcacgtacaggctcttgccctcctgcttggttcacgtagtagcagggtgttgcccacggacac  
ggtgtccacgcccttggttgacacgtagtc 3'
- 3)olfhnnum21.seq, bases 1413-1497 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mARN  
5'gagcctgtacgtgaagggcgagcccatcatcaacttctacgacccgctggtgttcccctccga  
cgagttcgacgcctccatctccc 3'
- 4)olfhnnum22.seq, bases 1483-1599 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary of mARN  
5'gttcatgatgttggtgggtggacttgccggcggttcacgttggtgcagcagctcgtcggacttgcg  
gatgaaggccaggctctggttgatcttctcgttcacctgggagatggaggcgtc 3'
- 5)olfhnnum23.seq, bases 1581-1691 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mARN  
5'caccaccaacatcatgaacaacgagttcatggaggtgaccgagaagatccagatggcctccga  
caacatcaacgacctgatccagtccggcggtgaacacccggctgctgac 3'
- 6)olfhnnum24.seq, bases 1677-1779 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary of mARN  
5'gatggtgatctcgctgatgaacttccgcaggtcggacatctgctgggtcagggagatggggat  
gtagttctgcacgtggtcttggtgatgggtcagcagccgggtg 3'
- 7)olfhnnum25.seq, bases 1761-1865 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'catcagcgagatcaccatccggaacgacaaccaggaggtgccccccagaggatcaccacga  
cgtgggcataaagcccctgaaccccgacgacttctggcgctg 3'
- 8)olfhnnum26.seq, bases 1849-1967 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary of mARN  
5'gtgcgcacgcagccgtccacggtggtgggcatggccagcaggccgggcccgggcatcagcctt  
atcttgggggtcttcatcagggaggggaggccggaggtgcagcgccagaagtcgtc 3'
- 9)olfhnnum27.seq, bases 1953-2059 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'cggctgctgctgcgcacccccctccctggtgatcaacgacctgatctacgcctacacctccaacct  
gatcaccgcggctgccaggacatcggcaagtcctaccaggtgc 3'
- 10)olfhnnum28.seq, bases 2043-2154 of F<sub>RSV</sub>HN<sub>Piv3</sub>, , inverse  
complementary of mARN  
5'ggacttcctggtgtcgttgatggtgaaggtgtgggagatccggggggttcaggtcgggcaccag  
gtcggagttcacggtgatgatgccgatctgcagcacctggtaggacttg

[SEQUENCES ABOVE ARE SEQ ID NOS: 30-39, respectively]

32/73

**Fig. 31: Sequence of the 16 oligonucleotides from which PCR fragment C was generated.**

- 1)olfhnum29.seq, bases 2139-2229 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'cgacaacaggaagtcctgctccctggccctcctgaacaccgacgtgtaccagctgtgctccac  
gcccaaggtggacgagcgctccgactac 3'
- 2)olfhnhum30.seq, bases 2214-2307 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mRNA  
5'gttcttgaagcgggtggtggagatggagccgtcgtggttgacgatgtccagcacgatgtcctc  
gatgccggagctggcgtagtcggagcgctcg 3'
- 3)olfhnhum31.seq, bases 2292-2398 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'caccgcgttcaagaacaacaacatcagcttcgaccagccctacgccgcctgtaccctccgt  
gggccccggcatctactacaagggaagatcatcttcctgggc 3'
- 4)olfhnhum32.seq, bases 2382-2472 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mRNA  
5'ccgctgggtcttgccggggcaccgggtggtggtgcagatggcgttctcgttgatgggggtgctc  
caggccgccgtagcccaggaagatgac 3'
- 5)olfhnhum33.seq, bases 2457-2549 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'cggcaagaccagcgggactgcaaccaggcctcccacagcccctgggttctccgaccgcccgc  
ggtgaactccatcatcgtggtggacaagg 3'
- 6)olfhnhum34.seq, bases 2532-2643 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mRNA  
5'cttggtgccagcagcagcaggcggccctcggagccccagtagttctgccgcatggagatggt  
ccacaccttcagcttggggatggagttcaggcccttgccaccacgatg 3'
- 7)olfhnhum35.seq, bases 2628-2726 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'gctgctgggcaacaagatctacatctacaccgctccaccagctggcacagcaagctgcagct  
gggcatcatcgacatcaccgactacagcgacatccg 3'
- 8)olfhnhum36.seq, bases 2710-2781 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mRNA  
5'ggggcactcgttggtgccggggccggctcagcacgttggtgccagggtccacttgatgcggatgtc  
gctgtagtc 3'
- 9)olfhnhum37.seq, bases 2765-2836 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'gcaacaacgagtggccctggggccactcctgccccgacggctgcatcaccggcgtgtacaccg  
acgcctacc 3'
- 10)olfhnhum38.seq, bases 2820-2889 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mRNA  
5'cttctgggagtcaggatcacggagctcacgatgctgccggtgggggttcagggggtaggcgtc  
ggtgtac

33/73

11)olfhnhum39.seq, bases 2874-2943 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'cctggactcccagaagtcccgggtgaaccccgtgatcacctacagcacctccaccgagcgcg  
gaacgag

12)olfhnhum40.seq, bases 2927-2994 of F<sub>RSV</sub>HN<sub>Piv3</sub>from: 1 to: 68,  
inverse complementary to mRNA  
5'gcagctggtggtggtgtagccggcgctcagggctcttggtgcggtatggccagctcgttcacgcg  
ctcgg

13)olfhnhum41.seq, bases 2979-3043 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'caccaccaccagctgcatcacccactacaacaagggctactgcttccacatcgtggagatcaa  
cc

14)olfhnhum42.seq, bases 3027-3085 of F<sub>RSV</sub>HN<sub>Piv3</sub>, inverse  
complementary to mRNA  
5'cggtcttgaacagcatgggctggaaggtgtccaggctcttggtggttgatctccacgatg 3'


15)olfhnhum43.seq, bases 3069-3114 of F<sub>RSV</sub>HN<sub>Piv3</sub>, homologous to mRNA  
5'catgctgttcaagaccgagatccccaagagctgcagctaaGAATTC 3'

[SEQUENCES ABOVE ARE SEQ ID NOS : 40-54, respectively]


34/73

**Fig. 32 : Construction of pEE14<sub>FRSVhum</sub>HN<sub>PIV3</sub>hum**



a) pNIV4120 +PCR fragment A



b) PCR fragment B



c) PCR fragment C

d) pEE14 F<sub>RSV</sub>humHN<sub>PIV3</sub>hum

35/73

**Fig. 33A : Humanized nucleic acids sequence of F<sub>HSV</sub>HN<sub>Piv3</sub> (upper sequence) compared to the original sequence found in the pNIV4120.**

```

7  AGAGGATCC.....ACCATGGAGCTGCTGATttttaAAGACCAACGCCA 49
   |||||
2262 AGAGGATCCCCCGGGTAccatggagttgctaatcctcaaaacaaatgcaa 2311

50  TCACCGCCATCCTGGCCGCGGTGACCCTCTGCTTCGCGTCCAGCCAGAAC 99
   | ||||| .||| | | | | | | | | | | | | | | | | | | |
2312 ttaccgcaatccttgctgcagtcacactctgttttgcttccagtcaaaac 2361

100 ATCACCGAGGAGTTCTACCAGTCCACCTGCTcttGCTGTCAGCAAGGGCTA 149
   ||||| | | | | | | | | | | | | | | | | | | | | |
2362 atcactgaagaattttatcaatcaacatgcagtgagtcagcaaaaggcta 2411

150 CCTGAGCGCGCTGAGGACGGGGTGGTACACTAGtGTGATCACCATCGAGC 199
   || | | | | | | | | | | | | | | | | | | | | |
2412 tcttagtgctctaagaactggttggtatactagtgttataactatagaat 2461

200 TGAGCAACATCAAGGAGAACAAGTGCAACGGCACCACGCAAGGTGAAG 249
   | | | | | | | | | | | | | | | | | | | | | | |
2462 taagtaatatcaaggaaaataagtgtaatggaacagacgctaaggtaaaa 2511

250 CTGATCAAGCAGGAGCTGGACAAGTACAAGAGCGCCGTGACCGAGCTGCA 299
   |||| | | | | | | | | | | | | | | | | | | | |
2512 ttgataaaaacaagaattagataaatataaaagtgtgtaacagaattgca 2561

300 GCTGCTGATGCAGAGCACCCCCGCCACCAACAACagaGCCAGGCGCGAGC 349
   | |||| | | | | | | | | | | | | | | | | | | |
2562 gttgctcatgcaaagcacaccggcaaccaacaatcgagccagaagagaac 2611

350 TGCCCAGGTTCATGAACTACACCCTCAACAACACCAAGAACACCAACGTG 399
   | | | | | | | | | | | | | | | | | | | | | | |
2612 taccaaggtttatgaattatacactcaacaataccaaaaataccaatgta 2661

400 ACCCTGAGCAAGAAGcGcAAGaggCGcTTCCTGGGCTTCCTGCTGGGCGT 449
   || | | | | | | | | | | | | | | | | | | | | |
2662 acattaagcaagaaaaggaaaagaagatttcttggtttttgtaggtgt 2711

450 GGGCTCCGCCATCGCCAGCGGCATCGCGGTGTCCAAGGTCCTGCACCTGG 499
   || | | | | | | | | | | | | | | | | | | | | |
2712 tggatctgcaatcgccagtggtcattgctgtatctaaggctcctgcacctag 2761

500 AGGGGGAGGTGAACAAGATCAAGAGCGCCCTGCTCTCCACCAACAAGGCG 549
   | |||| | | | | | | | | | | | | | | | | | | |
2762 aaggggaagtgaacaaaatcaaaagtgtctactatccacaaacaaggct 2811

```

550 GTGGTCAGCCTGTCCAACGGCGTGAGCGTGCTGACCAGCAAGGTGCTGGA 599  
|| ||| | | | | | | | | | | | | | |  
2812 gtagtcagcttatcaaatggagttagtgtcctaaccagcaaagtgttaga 2861

600 CCTCAAGAACTACATCGACAAGCAatTGCTCCCCATCGTGAACAAGCAGt 649  
||| | | | | | | | | | | | | | | | |  
2862 cctcaaaaactatatagataaacagttggttacctattgtgaacaagcaaa 2911

650 cCTGCAGCATCTCTAACATTGAGACCGTGATCGAGTTCCAGCAGAAGAAC 699  
|| | | | | | | | | | | | | | | | | | |  
2912 gctgtagcatatcaaacattgaaactgtgatagagttccaacaaaagaac 2961

700 AACAGGCTGCTGGAGATCAC CAGGGAGTTCAGCGTGAACGCgGGcGTcAC 749  
||| | | | | | | | | | | | | | | | |  
2962 aacagactactagagattaccagggaatttagtgtaaagcgaggtgtaac 3011

750 CACCCCGGTGAGCACCTACATGCTGACCAACAGCGAGCTGCTGTCCCTGA 799  
|| | | | | | | | | | | | | | | | | |  
3012 tacacctgtaagcacttatatgttaacaaatagtgaattattatcattaa 3061

800 TCAACGACATGCCCATCACCAACGACCAGAAGAAGCTtATGTCCAACAAC 849  
||| | | | | | | | | | | | | | | | |  
3062 tcaatgatatgcctataacaaatgatcagaaaaagttaatgtccaacaat 3111

850 GTGCAGATCGTGCGCCAGCAGAGCTACagCATCATGagCATCATCAAGGA 899  
|| | | | | | | | | | | | | | | | | |  
3112 gttcaaatagtttagacagcaaagttactctatcatgtccataataaagga 3161

900 GGAGGTGCTGGCCTACGTGGTGCAGCTGCCCCTGTACGGCGTGATCGACA 949  
|| | | | | | | | | | | | | | | | | |  
3162 ggaagtcttagcatatgtagtacaattaccactatatggtgtaatagata 3211

950 CCCCCTGCTGGAAGCTGCACACCTCCCCCCTGTGCACCACCAACACCAAG 999  
| | | | | | | | | | | | | | | | | |  
3212 caccttggttgaaactgcacacatcccctctatgtacaaccaacacaaag 3261

1000 GAGGGCTCCAACATCTGCCTGACCCGCACCGACCGGGGCTGGTACTGCGA 1049  
|| | | | | | | | | | | | | | | | | |  
3262 gaaggggtccaacatctgtttaacaagaaccgacagaggatggtactgtga 3311

1050 CAACGCCGGCTCCGTGTCCTTCTTCCCCCTGGCGGAGACCTGCAAGGTGC 1099  
|| | | | | | | | | | | | | | | | | |  
3312 caatgcaggatcagtatctttcttcccactagctgaaacatgtaaagttc 3361

1100 AGTCCAACCGCGTGTTCTGCGACACCATGAACAGCCTGACCCTGCCCAGC 1149  
| | | | | | | | | | | | | | | | | |

3362	aatcgaatcgagtattttgtgacacaatgaacagtttaacattaccaagt	3411
1150	GAGGTGAACCTCTGCAACATCGACATCTTCAACCCCAAGTACGACTGCAA	1199
3412	gaagtaaattctctgcaacattgacatattcaaccccaaataatgattgcaa	3461
1200	GATtATGacctccaagaccgacgtgagcagctccgtgatcacctccctgg	1249
3462	aattatgacttcaaaaacagatgtaagcagctccgttatcacatctctag	3511
1250	gcgccatcgtgtcctgctacggcaagaccaagtgtacagcctccaacaag	1299
3512	gagccattgtgtcatgctatggcaaaactaaatgtacagcatccaataaa	3561
1300	aaccgcggcatcatcaagaccttctccaacggctgcgactacgtgtccaa	1349
3562	aatcgtggaatcataaagacatttttctaacgggtgtgattatgtatcaaa	3611
1350	caagggcggtggacaccgtgtccgtgggcaacaccctgtactacgtgaaca	1399
3612	taaggggggtggacactgtgtctgtaggtataacattatattatgtaaata	3661
1400	agcaggagggcaagagcctgtacgtgaagggcgagcccatcatcaacttc	1449
3662	agcaagaaggcaaaaagtctctatgtaaaagggtgaaccaataataaatttc	3711
1450	tacgacccgctgggtgttcccctccgacgagttcgacgcctccatctccca	1499
3712	tatgaccattagtggttcccctctgatgaatttgatgcatcaatatctca	3761
1500	ggtgaacgagaagatcaaccagagcctggccttcacccgcaagtccgacg	1549
3762	agtcaatgagaagattaaccagagcctagcatttattcgtaaataccgatg	3811
1550	agctgctgcacaacgtgaacgccggcaagtccaccaccaacatcatgaac	1599
3812	aattattacataatgtaaatactgtggtaaatccaccacaaatatcatgAAC	3861
1600	aacgagttcatggaggtgaccgagaagatccagatggcctccgacaacat	1649
3862	AATGAGTTTATGGAAGTTACAGAAAAGATCCAAATGGCATCGGATAATAT	3911
1650	caacgacctgatccagtcggcggtgaacacccggctgctgaccatccaga	1699
3912	TAATGATCTAATACAGTCAGGAGTGAATACAAGGCTTCTTACAATTTCAGA	3961

38/73

1700 gccacgtgcagaactacatccccatctccctgacccagcagatgtccgac 1749  
| | | | | | | | | | | | | | | | | | | | | |  
3962 GTCATGTCCAGAATTATATACC<sup>a</sup>ATATCATTGACACAACAAATGTCGGAT 4011  
  
1750 ctgcggaagttcatcagcgagatcaccatccggaacgacaaccaggaggt 1799  
| | | | | | | | | | | | | | | | | | | | | |  
4012 CTTAGGAAATTCATTAGTGAAATTACAATTAGGAATGATAATCAAGAAGT 4061  
  
1800 gccccccagaggatcacccacgacgtgggcataaagcccctgaaccccg 1849  
| | | | | | | | | | | | | | | | | | | | | |  
4062 GCCTCCACAAAGAATAACACATGATGTGGGCATAAAACCTTTAAATCCAG 4111  
  
1850 acgacttctggcgctgcacctccggcctccccctccctgatgaagaccccc 1899  
| | | | | | | | | | | | | | | | | | | | | |  
4112 ATGATTTTTGGAGATGCACGTCTGGTCTTCCATCTTTAATGAAAACCTCCA 4161  
  
1900 aagataaggctgatgcccgggcccgccctgctggccatgccaccaccgt 1949  
| | | | | | | | | | | | | | | | | | | | | |  
4162 AAAATAAGGTTAATGCCGGGGCCGGGATTATTAGCTATGCCAACGACTGT 4211  
  
1950 ggacggctgcgtgcgcacccccctccctgggtgatcaacgacctgatctacg 1999  
| | | | | | | | | | | | | | | | | | | | | |  
4212 TGATGGCTGTGTTAGAACTCCGTCCTTAGTTATAAATGATCTGATTTATG 4261  
  
2000 cctacacctccaacctgatcacccgcggctgccaggacatcggcaagtcc 2049  
| | | | | | | | | | | | | | | | | | | | | |  
4262 CTTATACCTCGAATCTAATTACTCGAGGTTGCCAGGATATAGGAAAATCA 4311  
  
2050 taccagggtgctgcagatcggcatcatcacctgaactccgacctggtacc 2099  
| | | | | | | | | | | | | | | | | | | | | |  
4312 TATCAAGTATTACAGATAGGGATAATAACTGTAAACTCAGACTTGGTACC 4361  
  
2100 cgacctgaacccccggatctcccacaccttcaacatcaacgacaacagga 2149  
| | | | | | | | | | | | | | | | | | | | | |  
4362 TGACTTAAATCCTAGGATCTCTCATACTTTCAACATAAATGACAATAGAA 4411  
  
2150 agtcctgctccctggccctcctgaacaccgacgtgtaccagctgtgctcc 2199  
| | | | | | | | | | | | | | | | | | | | | |  
4412 AGTCATGTTCTCTAGCACTCCTAAACACAGATGTATATCAACTGTGTTCG 4461  
  
2200 acgcccgaaggtggacgagcgctccgactacgccagctccggcatcgagga 2249  
| | | | | | | | | | | | | | | | | | | | | |  
4462 ACTCCCAAAGTTGATGAAAGATCAGATTATGCATCATCAGGCATAGAAGA 4511  
  
2250 catcgtgctggacatcgtcaaccacgacggctccatctccaccacccgct 2299  
| | | | | | | | | | | | | | | | | | | | | |  
4512 TATTGTACTTGATATtGTCAATCATGATGGTTCAATCTCAACAACAAGAT 4561

[illegible]

DEC 04 2000 18:52

41/73

Fig. 33B

(Linear) MAP of: FrsvHNpiv3.seq check: 7448 from: 1 to:  
3090 nucleic acids sequence of FrsvHNpiv3 (non humanised)

atggagttgctaatacctcaaaacaaatgcaattaccgcaatccttgctgc  
agtcacactctgttttgcttccagtcaaaacatcactgaagaattttatc  
aatcaacatgcagtgagtcagcaaaggctatcttagtgctctaagaact  
ggttggtatactagtggtataactatagaattaagtaatatcaaggaaaa  
taagtgtaatggaacagacgctaaggtaaaattgataaaacaagaattag  
ataaatataaaaagtgctgtaacagaattgcagttgctcatgcaaagcaca  
ccggcaaccaacaatcgagccagaagagaactaccaagggtttatgaatta  
tacactcaacaataccaaaaataccaatgtaacattaagcaagaaaagga  
aaagaagatttcttggttttttggttaggtggtggatctgcaatcgccagt  
ggcattgctgtatctaagggtcctgcacctagaaggggaagtgaacaaaat  
caaaagtgtctactatccacaaacaaggctgtagtcagcttatcaaatg  
gagttagtggtcttaaccagcaaagtgttagacctcaaaaactatatagat  
aaacagttgttacctattgtgaacaagcaaagctgtagcatatcaaacat  
tgaaactgtgatagagttccaacaaaagaacaacagactactagagatta  
ccaggggaatttagtggttaatgcaggtgtaactacacctgtaagcacttat  
atgttaacaaatagtggaattattatcattaatcaatgatatgcctataac  
aatgatcagaaaaagttaatgtccaacaatgttcaaatagttagacagc  
aaagttactctatcatgtccataataaaggaggaagtcttagcatatgta  
gtacaattaccactatatggtgtaatagatacaccttggttggaactgca  
cacatcccctctatgtacaaccaacacaaaggaaggggtccaacatctgtt  
taacaagaaccgacagaggatggtactgtgacaatgcaggatcagtatct

42/73

ttcttcccactagctgaaacatgtaaagttcaatcgaatcgagtattttg  
tgacacaatgaacagtttaacattaccaagtgaagtaaattctctgcaaca  
ttgacatattcaacccccaaatatgattgcaaaattatgacttcaaaaaca  
gatgtaagcagctccggttatcacatctctaggagccattgtgtcatgcta  
tggcaaaaactaaatgtacagcatccaataaaaaatcgtggaatcataaaga  
catttttctaacgggtgtgattatgtatcaaataaggggggtggacactgtg  
tctgtaggtaatacattatattatgtaaataagcaagaaggcaaaagtct  
ctatgtaaaaggtgaaccaataataaatttctatgaccatttagtggtcc  
cctctgatgaatttgatgcatcaatatctcaagtcaatgagaagattaac  
cagagcctagcattttattcgtaaatccgatgaattattacataatgtaa  
tgctggtaaatccaccacaaatatcatgAACAATGAGTTTATGGAAGTTA  
CAGAAAAGATCCAAATGGCATCGGATAATATTAATGATCTAATACAGTCA  
GGAGTGAATACAAGGCTTCTTACAATTCAGAGTCATGTCCAGAATTATAT  
ACCaATATCATTGACACAACAAATGTCGGATCTTAGGAAATTCATTAGTG  
AAATTACAATTAGGAATGATAATCAAGAAGTGCCTCCACAAAGAATAACA  
CATGATGTGGGCATAAAACCTTTAAATCCAGATGATTTTTGGAGATGCAC  
GTCTGGTCTTCCATCTTTAATGAAAACCTCCAAAATAAGGTTAATGCCGG  
GGCCGGGATTATTAGCTATGCCAACGACTGTTGATGGCTGTGTTAGAACT  
CCGTCCTTAGTTATAAATGATCTGATTTATGCTTATACCTCaAATCTAAT  
TACTCGAGGTTGCCAGGATATAGGAAAATCATATCAAGTATTACAGATAG  
GGATAATAACTGTAAACTCAGACTTGGTACCTGACTTAAATCCTAGGATC  
TCTCATACTTTCAACATAAATGACAATAGAAAGTCATGTTCTCTAGCACT  
CCTAAAtACAGATGTATATCAACTGTGTTCGACTCCCAAAGTTGATGAAA  
GATCAGATTATGCATCATCAGGCATAGAAGATATTGTACTTGATATtGTC

43/73

AATCATGATGGTTCAATCTCAACAACAAGATTTAAGAACAATAATATAAG  
TTTTGATCAACCATATGCGGCATTATACCCATCTGTTGGACCAGGGATAT  
ACTACAAAGGCAAAATAATATTTCTCGGGTATGGAGGTCTTGAACATCCA  
ATAAATGAGAATGCAATCTGCAACACAACCTGGGTGTCCCGGGAAAACGCA  
GAGAGACTGCAATCAGGCATCTCATAGTCCcTGGTTTTTCAGACAGAAGGA  
TGGTCAACTCCATTATTGTTGTTGACAAGGGCTTAAACTCAATTCCAAAA  
CTGAAGGTATGGACGATATCCATGAGACAAAATTACTGGGGGTCAGAAGG  
AAGGCTACTTCTACTAGGTAACAAGATCTATATATATACAAGATCTACAA  
GTTGGCATAGCAAGTTACAATTAGGAATAATTGATATTACTGATTACAGT  
GATATAAGAATAAAATGGACATGGCATAATGTGtTATCAAGACCAGGAAA  
CAATGAATGTCCATGGGGACATTCATGtCCAGATGGATGTATAACAGGAG  
TATATACTGATGCATATCCgCTCAATCCCACAGGGAGCATTGTGTCATCT  
GTCATATTAGACTCGCAAAAATCGAGAGTAAACCCAGTCATAACTTACTC  
AACAtCAACTGAAAGGGTAAACGAGCTGGCCATCCGAAACAAAACACTCT  
CAGCTGGATATAACAACGAGCTGCATTACACACTATAACAAAGGATAT  
TGTTTTTCATATAGTAGAAATAAATCATAAAAGCTTAGACACATTCCAACC  
TATGTTGTTCAAAACAGAGATTCCAAAAAGCTGCAGTTAA

[SEQ ID NO: 55]

44/73

Fig. 33C

(Linear) MAP of: FrhumHNphum.seq check: 9920 from: 1 to:  
3090 Humanised nucleic acids sequence of FRSVHNPIV3

GGGTGGTACACTAGtGTGATCACCATCGAGCTGAGCAACATCAAGGAGAA  
CAAGTGCAACGGCACCGACGCCAAGGTGAAGCTGATCAAGCAGGAGCTGG  
ACAAGTACAAGAGCGCCGTGACCGAGCTGCAGCTGCTGATGCAGAGCACC  
CCCGCCACCAACAACagaGCCAGGCGCGAGCTGCCCAGGTTTCATGAACTA  
CACCTCAACAACACCAAGAACACCAACGTGACCCTGAGCAAGAAGcGcA  
AGaggCGcTTCCTGGGCTTCCTGCTGGGCGTGGGCTCCGCCATCGCCAGC  
GGCATCGCGGTGTCCAAGGTCCTGCACCTGGAGGGGGAGGTGAACAAGAT  
CAAGAGCGCCCTGCTCTCCACCAACAAGGCGGTGGTCAGCCTGTCCAACG  
GCGTGAGCGTGCTGACCAGCAAGGTGCTGGACCTCAAGAACTACATCGAC  
AAGCAatTGCTCCCCATCGTGAACAAGCAGtcCTGCAGCATCTCTAACAT  
TGAGACCGTGATCGAGTTCCAGCAGAAGAACAACAGGCTGCTGGAGATCA  
CCAGGGAGTTCAGCGTGAACGCgGGcGTcACCACCCCGGTGAGCACCTAC  
ATGCTGACCAACAGCGAGCTGCTGTCCCTGATCAACGACATGCCCCATCAC  
CAACGACCAGAAGAAGCTtATGTCCAACAACGTGCAGATCGTGCGCCAGC  
AGAGCTACagCATCATGagCATCATCAAGGAGGAGGTGCTGGCCTACGTG  
GTGCAGCTGCCCCCTGTACGGCGTGATCGACACCCCCTGCTGGAAGCTGCA  
CACCTCCCCCCTGTGCACCACCAACACCAAGGAGGGCTCCAACATCTGCC  
TGACCCGCACCGACCGGGGCTGGTACTGCGACAACGCCGGCTCCGTGTCC  
TTCTTCCCCCTGGCGGAGACCTGCAAGGTGCAGTCCAACCGCGTGTTCTG  
CGACACCATGAACAGCCTGACCCTGCCCAGCGAGGTGAACCTCTGCAACA

45/73

TCGACATCTTCAACCCCAAGTACGACTGCAAGATtATGacctccaagacc  
gacgtgagcagctccgtgatcacctccctgggcgccatcgtgtcctgcta  
cggcaagaccaagtgtacagcctccaacaagaaccgcggcatcatcaaga  
ccttctccaacggctgcgactacgtgtccaacaagggcgtggacaccgtg  
tccgtgggcaacaccctgtactacgtgaacaagcaggagggcaagagcct  
gtacgtgaagggcgagcccatcatcaacttctacgaccgcgtggtgttcc  
cctccgacgagttcgacgcctccatctcccagggtgaacgagaagatcaac  
cagagcctggccttcatccgcaagtccgaacgagctgctgcacaacgtgaa  
cgccggcaagtccaccaccaacatcatgaacaacgagttcatggagggtga  
ccgagaagatccagatggcctccgacaacatcaacgacctgatccagtcc  
ggcgtgaacaccggctgctgaccatccagagccacgtgcagaactacat  
ccccatctccctgaccagcagatgtccgacctgcggaagttcatcagcg  
agatcaccatccggaacgacaaccaggaggtgccccccagaggatcacc  
cacgacgtgggcataaagcccctgaaccccgacgacttctggcgctgcac  
ctccggcctcccctccctgatgaagaccccccaagataaggctgatgcccg  
ggcccggcctgctggccatgcccaccaccgtggacggctgcgtgcgcacc  
ccctccctggtgatcaacgacctgatctacgcctacacctccaacctgat  
caccgcggctgccaggacatcggcaagtcctaccagggtgctgcagatcg  
gcatcatcaccgtgaactccgacctggtaccgcacctgaacccccggatc  
tcccacaccttcaacatcaacgacaacaggaagtcctgctccctggccct  
cctgaacaccgacgtgtaccagctgtgctccacgcccagggtggacgagc  
gctccgactacgccagctccggcatcgaggacatcgtgctggacatcgtc  
aaccacgacggctccatctccaccaccgcgttcaagaacaacaacatcag  
cttcgaccagccctacgccgcctgtaccctccgtgggccccggcatct


46/73

actacaagggcaagatcatcttcctgggctacggcggcctggagcacccc  
atcaacgagaacgccatctgcaacaccaccgggtgccccggcaagacca  
gcgggactgcaaccaggcctcccacagcccctggttctccgaccgccgca  
tggtgaactccatcatcgtgggtggacaagggcctgaactccatccccaag  
ctgaagggtgtggaccatctccatgcggcagaactactggggctccgaggg  
ccgcctgctgctgctgggcaacaagatctacatctacacccgctccacca  
gctggcacagcaagctgcagctgggcatcatcgacatcacggactacagc  
gacatccgcatcaagtggacctggcacaacgtgctgagccgggccggcaa  
caacgagtgcccctggggccactcctgccccgacggctgcatcacggcg  
tgtacaccgacgcctaccccctgaaccccaccggcagcatcgtgagctcc  
gtgatcctggactcccagaagtcccgggtgaaccccgtgatcacctacag  
cacctccaccgagcgcgtgaacgagctggccatccgcaacaagacctga  
gcgccggctacaccaccaccagctgcatcacccactacaacaagggctac  
tgcttccacatcgtggagatcaaccacaagagcctggacaccttccagcc  
catgctgttcaagaccgagatccccaagagctgcagctaa


[SEQ ID NO : 56]

47/73

Fig. 34A



48/73

Fig. 34B: Humanization impact on the level of expression of  $F_{RSV}HN_{FluV3}$ 

49/73


Fig 35: Codon usage of  $F_{Muv}H_{Mv}$  and highly expressed human genes (hum high exp)

The frequencies (x100) of the individual codons are shown for each of the degenerately encoded amino acids, and the most prevalent codon is shown in bold.

		hum high exp	$F_{Muv}H_{Mv}$			hum high exp	$F_{Muv}H_{Mv}$			hum high exp	$F_{Muv}H_{Mv}$			hum high exp	$F_{Muv}H_{Mv}$
Ala	GCG	17	10	Gln	CAG	<b>88</b>	32	Leu	TTG	6	<b>25</b>	Ser	AGT	10	19
	A	13	<b>47</b>		A	12	<b>68</b>		A	2	11		C	<b>34</b>	15
	T	17	25						CUG	<b>58</b>	15		TCG	9	5
	C	<b>53</b>	18						A	3	20		A	5	<b>30</b>
									T	5	14		T	13	17
									C	26	15		C	28	14
Arg	AGG	18	<b>30</b>	Glu	GAG	<b>75</b>	52					Thr	ACG	15	7
	A	10	<b>30</b>		A	25	<b>48</b>						A	14	<b>41</b>
	CGG	21	10										T	14	30
	A	6	15										C	<b>57</b>	21
	T	7	8	Gly	GGG	24	24	Lys	AAG	<b>82</b>	<b>51</b>				
	C	<b>37</b>	8		A	14	26		A	18	49				
					T	12	<b>33</b>								
					C	<b>50</b>	17								
Asn	AAT	22	<b>60</b>					Phe	TTT	20	39				
	C	<b>78</b>	40						C	<b>80</b>	<b>61</b>	Tyr	TAT	26	34
													C	<b>74</b>	<b>66</b>
Asp	GAT	25	<b>59</b>	His	CAT	21	<b>60</b>								
	C	<b>75</b>	41		C	<b>79</b>	40								
Cys	TGT	32	<b>56</b>	Ile	ATA	5	25	Pro	CCG	17	19	Val	GTG	<b>64</b>	<b>34</b>
	C	<b>68</b>	44		T	18	35		A	16	<b>40</b>		A	5	10
					C	<b>77</b>	<b>39</b>		T	19	23		T	7	27
									C	<b>48</b>	19		C	25	29

50/73

**Fig. 36: Schematic diagram of the PCR synthesis of each fragment in which X and Y are restriction sites that allow retrieval of the full size fragment from the cloning vector.**



51/73

**Fig 37 : Sequence of the 12 oligonucleotides from which PCR fragment A was generated.**

1) oli 1 FmuvHmv 1-98, hom ARN  
ggtctagaccaccATGAAGGCGTTCCCCGTGATCTGCCTGGGCTTCGCCATCTTCTCCAG  
-----+-----+-----+-----+-----+-----+ 60  
CAGCATCTGCGTGAACATCAACATCCTGCAGCAGATCG  
-----+-----+-----+-----+-----+ 98

2) oli 2 FmuvHmv 82-181, inv comp ARN  
GTTGGGCAGCAGCTTGACCACCACGTAGGAGCTGGAGCTCTGGGAGTAGTAGCTCAGCTG  
-----+-----+-----+-----+-----+-----+ 60  
CCTCACCTGCTGCTTGATGTATCCGATCTGCTGCAGGATG  
-----+-----+-----+-----+-----+ 100

3) oli 3 FmuvHmv, 166-264 hom ARN  
CAAGCTGCTGCCCAACATCCAGCCCACCGACAACAGCTGCGAGTTCAAGAGCGTGACCCA  
-----+-----+-----+-----+-----+-----+ 60  
GTACAACAAGACCCTGAGCAACCTGCTGCTGCCCATCGC  
-----+-----+-----+-----+-----+ 99

4) oli 4 FmuvHmv, 250-352, inv comp ARN  
CAGGGCGGCGATGCCGATGGCGATGCCGGCGAACCGCTTGTGCCGCGGGAGCCGGGGGA  
-----+-----+-----+-----+-----+-----+ 60  
GGGGGAGGTGATGTTGTTGATGTTCTCGGCGATGGGCAGCAGC  
-----+-----+-----+-----+-----+ 103

5) oli 5 FmuvHmv, 338-441, hom ARN  
GGCATCGCCGCCCTGGGCGTGGCCACCGCCGCCAGGTGACCGCCGCGTGTCCCTGGTG  
-----+-----+-----+-----+-----+-----+ 60  
CAGGCCCAGACCAACGCCCGCGCCATCGCCGCCATGAAGAACTC  
-----+-----+-----+-----+-----+ 104

6) oli 6 FMuvHmv, 427-523, inv comp ARN  
GTCCTGGATGGCCTGCACGGCGATGGCCAGCTGCTGGGTGCCCTCCTTCACCTCGAACAC  
-----+-----+-----+-----+-----+-----+ 60  
GGCGCGGTTGGTGGCCTGGATGGAGTTCTTCATGGCG  
-----+-----+-----+-----+-----+ 97

7) oli 7 FMUVHM 509-610, hom ARN  
CAGGCCATCCAGGACCACATCAACACCATCATGAACACCCAGCTGAACAACATGTCCTGC  
-----+-----+-----+-----+-----+-----+ 60  
CAGATCCTGGACAACCAGCTGGCCACCTCCCTGGGCCTGTAE  
-----+-----+-----+-----+-----+ 102

52/73

8) oli 8 FMUHM, 595-691, inv comp ARN  
GGACCGCAGGGCCTGGATACTGATGGGggaCAGGGCGGGGTTGATCAGCTGGGGCTGGAA 60  
-----+-----+-----+-----+-----+-----+-----+-----+  
CACGGTGGTCAGCTCGGTACAGGTACAGGCCAGGGAG 97  
-----+-----+-----+-----+-----+-----+-----+-----+

9) oli 9, 677-778, hom ARN  
CAGGCCCTGCGGTCCCTGCTGGGCAGCATGACCCCGCCGTGGTGCAGGCCACCCTGAGC 60  
-----+-----+-----+-----+-----+-----+-----+-----+  
ACCTCCATCAGCGCCGCCGAGATCCTGAGCGCCGGCCTGATG 102  
-----+-----+-----+-----+-----+-----+-----+-----+

10) oli 10, FmuvHmv, 763-862, inv comp ARN  
GTTGGACTGGGTCACGATGGTGGGCACGTTGATCTTCACGATCATCTGCATCTCGTCCAG 60  
-----+-----+-----+-----+-----+-----+-----+-----+  
CAGCACGGACACGATCTGGCCCTCCATCAGGCCGGCGCTC 100  
-----+-----+-----+-----+-----+-----+-----+-----+

11) oli 11, FmuvHmv, 848-949, homARN  
GTGACCCAGTCCAACGCCCTGGTGATCGACTTCTACAGCATCAGCAGCTTCATCAACAAC 60  
-----+-----+-----+-----+-----+-----+-----+-----+  
CAGGAGTCCATCATCCAGCTGCCCCACCGCATCCTGGAGATC 102  
-----+-----+-----+-----+-----+-----+-----+-----+

12) oli 12 FMUHM, 935-1039, inv compARN  
GCTCAGCCGCTCGGCCTCGTTGTACTGGCAGAAGATGTGGTGGCGGGTCAGCTTGCAGTT 60  
-----+-----+-----+-----+-----+-----+-----+-----+  
CTTGGCGGGGTAGCGCCACTGCTCGTTGCCGATCTCCAGGATGCG 105  
-----+-----+-----+-----+-----+-----+-----+-----+

[SEQ ID NOS: 57-68 respectively]

53/73

**Fig. 38 : Sequence of the 9 oligonucleotides from which PCR fragment B was generated.**

12) oli 12 FMUHM, 935-1039, inv compARN  
GCTCAGCCGCTCGGCCTCGTTGTACTGGCAGAAGATGTGGTGGCGGGTCAGCTTGCAGTT 60  
-----+-----+-----+-----+-----+-----+-----+  
CTTGGCGGGGTAGCGCCACTGCTCGTTGCCGATCTCCAGGATGCG 105  
-----+-----+-----+-----+-----+-----+-----+

13) oli 13, 1025-1129, hom ARN  
GCCGAGCGGCTGAGCCTGGAGACCAAGCTGTGCCTGGCCGGCAACATCAGCGCCTGCGTG 60  
-----+-----+-----+-----+-----+-----+-----+  
TTCTCCAGCATCGCCGGCAGCTACATGCGCCGCTTCGTGGCCCTG 105  
-----+-----+-----+-----+-----+-----+-----+

14) oli 14, 1115-1216, inv comp ARN  
GGCGTGGTGGTCTGGGCTGGTAGATGGGGTAGGAGGGGCTCTTGACAGGCAGGTCAGGCT 60  
-----+-----+-----+-----+-----+-----+-----+  
GCGGCAGTTGGCCACGATGGTGCCGTCCAGGGCCACGAAGCG 102  
-----+-----+-----+-----+-----+-----+-----+

15) oli 15, 1202-1299, hom ARN  
CCCGACCACCACGCCGTGACCACCATCGACCTGACCTCCTGCCAGACCCTGAGCCTGGAC 60  
-----+-----+-----+-----+-----+-----+-----+  
GGCCTGGACTTCAGCATCGTGTCCCTGAGCAACATCAC 98  
-----+-----+-----+-----+-----+-----+-----+

16) oli 16 1285-1387, inv comp ARN  
CTTGCTCAGCTCGGTGGAGATGTCGATGGGCTGGGTGTTGATGGTCTGGCTCAGGCTGAT 60  
-----+-----+-----+-----+-----+-----+-----+  
GGTCAGGTTCTCGGCGTAGGTGATGTTGCTCAGG 94  
-----+-----+-----+-----+-----+-----+-----+

17) oli 17, 1363-1462, hom /ARN  
CACCGAGCTGAGCAAGGTGAACGCCTCCCTGCAGAACGCCGTGAAGTACATCAAGGAGAG 60  
-----+-----+-----+-----+-----+-----+-----+  
CAACCACCAGCTGCAGAGCGTGAGCGTGAGCAGCAAGCGC 100  
-----+-----+-----+-----+-----+-----+-----+

18) oli 18, 1447-1550, inv comp ARN  
TCACCTGGTGCTCGATGGAGTTGGTCACGTCCAGGTTGGTGCTCAGGCTCTTGTGGATCT 60  
-----+-----+-----+-----+-----+-----+-----+  
CGGCGGTGTAGATGGCGGCGCGGTGCAGGCGCTTGCTGCTCAG 104  
-----+-----+-----+-----+-----+-----+-----+

54/73

19) oli 19, 1534-1636, hom ARN  
CATCGAGCACCAGGTGAAGGACGTGCTGACCCCCCTGTTCAAGATCATCGGCGACGAGGT  
-----+-----+-----+-----+-----+-----+ 60  
GGGCCTGCGCACCCCCCAGCGCTTCACCGACCTGGTGAAGTTC  
-----+-----+-----+-----+-----+ 103

20) oli 20 FmuvHmv, 1622-1718, inv comp ARN  
GCTCGGGGGGGTTGATGCACCAGGTCAGGTCGCGGAAGTCGTACTCGCGGTCGGGGTTCA  
-----+-----+-----+-----+-----+-----+ 60  
GGAACTTGATCTTGTCGGAGATGAACTTCACCAGGTC  
-----+-----+-----+-----+-----+ 97

[SEQ ID NOS: 69-77 respectively]

55/73

**Fig. 39 : Sequence of the 11 oligonucleotides from which PCR fragment C was generated.**

20) oli 20 FmuvHmv, 1622-1718, inv comp ARN  
GCTCGGGGGGGTTGATGCACCAGGTCAGGTCGCGGAAGTCGTA CTCTCGCGGTCGGGGTTCA  
-----+-----+-----+-----+-----+-----+ 60  
GGA ACTTGATCTTGTCGGAGATGAACTTCACCAGGTC  
-----+-----+-----+-----+-----+ 97

21) oli 21, FmuvHmv, 1701-1799, hom ARN  
GCATCAACCCCCCGAGCGGATCAAGCTGGACTACGACCAGTACTGCGCCGACGTGGCCG  
-----+-----+-----+-----+-----+-----+ 60  
CCGAGGAGCTGATGAACGCCCTGGTGAACAGCACCCCTGC  
-----+-----+-----+-----+-----+ 99

22) oli 22, 1784-1888, inv comp  
CATGTTGCTGAACTGGCCCCGGATGGTGGTGGGGCCGCTGCAGTTGCCCTTGCTCACGGC  
-----+-----+-----+-----+-----+-----+ 60  
CAGGA ACTGGTTGGTGGTGC GGGTCTCCAGCAGGGTGCTGTTCAC  
-----+-----+-----+-----+-----+ 105

23) oli 23, 1874-1971, hom ARN  
CAGTTCAGCAACATGAGCCTGTCCCTGCTGGACCTGTACCTGGGCCGGGGCTACAACGTG  
-----+-----+-----+-----+-----+-----+ 60  
AGCAGCATCGTGACCATGACCAGCCAGGGCATGTACGG  
-----+-----+-----+-----+-----+ 98

24) oli 24, 1957-2057, inv.comp ARN  
CCACCTCGAACACGCGGTACATGCTCAGCTGGCTCAGCTCGCTCCGCTTGCTGCTCAGGT  
-----+-----+-----+-----+-----+-----+ 60  
TGGGCTTCTCCACCAGGTAGGTGCCGCCGTACATGCCCTGG  
-----+-----+-----+-----+-----+ 101

25) oli 25, FmuvHmv, 2043-2140, homARN  
GCGTGTTTCGAGGTGGGCGTGATCCGGAACCCCGCCTGGGCGCCCCCGTGTTCCACATGA  
-----+-----+-----+-----+-----+-----+ 60  
CCA ACTACCTGGAGCAGCCCGTGAGCAACGACCTGAGC  
-----+-----+-----+-----+-----+ 98

26) oli26, FmuvHmv, 2125-2227, inv compARN  
GCCGCTGCCCTGGTAGGGGATGGTGTGCTGTCCTCGCCGTGGCACAGGGCGGCCAGCTT  
-----+-----+-----+-----+-----+-----+ 60  
CAGCTCGCCCAGGGCCACCATGCAGTTGCTCAGGTCGTTGCTC  
-----+-----+-----+-----+-----+ 103

27) oli 27, 2212-2309, FmuvHm, hom ARN  
CTACCAGGGCAGCGGCAAGGGCGTGAGCTTCCAGCTGGTGAAGCTGGGCGTGTGGAAGAG  
-----+-----+-----+-----+-----+-----+ 60  
CCCCACCGACATGCAGAGCTGGGTGCCCTGAGCACCG  
-----+-----+-----+-----+-----+ 98

56/73

28) oli 28, FmuvHmv, 2294-2392, inv comp ARN  
GGTGGGCACGGCCCACTTGGCCTGGTTGTCGGCGATCACGCCGCGGTGGCTGCTCAGGTA 60  
-----+-----+-----+-----+-----+-----+  
CAGGCGGTTCGATCACGGGGTCGTCGGTGCTCAGGGGCAC 99  
-----+-----+-----+-----+-----+-----+

29) oli 29, Fmuv Hmv, 2377-2477, hom ARN  
GTGGGCCGTGCCCACCAACCCGCACCGACGACAAGCTGCGCATGGAGACCTGCTTCCAGCA 60  
-----+-----+-----+-----+-----+-----+  
GGCCTGCAAGGGCAAGATCCAGGCCCTGTGCGAGAACCCCG 101  
-----+-----+-----+-----+-----+-----+

30) oli 30, FmuvHmv 2462-2561, inv comp  
TGATCTTCAGCTCCACGGTCAGGCTCAGGTCCACGCTCAGCACGCCGTAGCTGGGGATGC 60  
-----+-----+-----+-----+-----+-----+  
GGTTGTCCTTCAGGGGGGGCCCAATCGGGGTTCTCGCACAG 100  
-----+-----+-----+-----+-----+-----+

[SEQ ID NOS: 78-88 respectively]

57/73

**Fig. 40 : Sequence of the 8 oligonucleotides from which PCR fragment D was generated.**

30) oli 30, FmuvHmv 2462-2561, inv comp  
TGATCTTCAGCTCCACGGTCAGGCTCAGGTCACGCTCAGCACGCCGTAGCTGGGGATGC  
-----+-----+-----+-----+-----+-----+ 60  
GGTTGTCCTTCAGGGGGGGCCCA~~t~~TCGGGGTTCTCGCACAG  
-----+-----+-----+-----+-----+ 100

31) oli31, FmuvHmv, 2546-2649, hom ARN  
GTGGAGCTGAAGATCAAGATCGCGAGCGGCTTCGGCCCCCTGATCACCCACGGCAGCGGC  
-----+-----+-----+-----+-----+-----+ 60  
ATGGACCTGTACAAGAGCAACCACAACAACGTGTACTGGCTGAC  
-----+-----+-----+-----+-----+ 104

32) oli 32, FmuvHmv, 2635-2738, inv comp ARN  
CGGTGAACAGGTAGGGGCTCACCTTGAAGCGGGGaATCCACTCCAGGGTGTTGATCACGC  
-----+-----+-----+-----+-----+-----+ 60  
CCAGGGCCAGGTTCTTCATGGGGGGGATGGTCAGCCAGTACACG  
-----+-----+-----+-----+-----+ 104

33) oli 33, FmuvHmv, 2723-2827, hom ARN  
CCCTACCTGTTCACCGTGCCCATCAAGGAGGCCGGCGAGGACTGCCACGCCCCGACCTAC  
-----+-----+-----+-----+-----+-----+ 60  
CTGCCCCGCCGAGGTGGACGGCGACGTGAAGCTGAGCAGCAACCTG  
-----+-----+-----+-----+-----+ 105

34) oli 34, FMUVHmv, 2813-2911, inv comp ARN  
CACGTAGTACACCACGGCGTGCTCCACGCGGCTGGTGTCTAGGTGGCCAGCACGTACTG  
-----+-----+-----+-----+-----+-----+ 60  
CAGGTCCTGGCCGGGCAGGATCACCAGGTTGCTGCTCAG  
-----+-----+-----+-----+-----+ 99

35) oli 35 FMUHM, 2897-2995, homARN  
GTGGTGTACTACGTGTACAGCCCCGGCCGAGCTTCTTCTACTTCTACCCCTTCCGCCTG  
-----+-----+-----+-----+-----+-----+ 60  
CCCATCAAGGGCGTGCCCATCGAGCTGCAGGTGGAGTGC  
-----+-----+-----+-----+-----+ 99

36) oli 36, FmuvHmv, 2981-3078, inv comp ARN  
CCGCTGTGGGTGATGTGGCCGCCGCTCTCGCTGTCGGCCAGCACGCAGAAGTGGCGGCAC  
-----+-----+-----+-----+-----+-----+ 60  
CACAGCTTCTGGTCCCAGGTGAAGCACTCCACCTGCAG  
-----+-----+-----+-----+-----+ 98

58/73

37) oli 37, 3064-3147, homARN

CATCACCCACAGCGGCATGGTGGGCATGGGCGTGAGCTGCACCGTGACCCGCGAGGACGG

-----+-----+-----+-----+-----+-----+ 60

CACCAACCGCCGCTAGcgaattcc


-----+-----+----- 84

[SEQ ID NOS: 89-96 respectively]


59/73

**Fig. 41 : Construction of pEE14F<sub>MuV</sub>hum HN<sub>MV</sub>hum**


a) PCR fragment A





b) pCR fragment B



c) PCR fragment C



d) PCR fragment D

d) pEE14 F<sub>MuV</sub>hum H<sub>MV</sub>hum

3

DEC 04 2000 18:57

1214	GCCGTGACCACCATCGACCTGACCTCCTGCCAGACCCTGAGCCTGGACGG	1263
1201	GCAGTCACGACCATTGATCTAACGTCATGTCAAACATTGTCCTGGACGG	1250
1264	CCTGGACTTCAGCATCGTGTCCCTGAGCAACATCACCTACGCCGAGAACC	1313
1251	ACTGGATTTTCAGCATTGTCTCGCTAAGCAACATCACTTACGCTGAGAATC	1300
1314	TGACCATCAGCCTGAGCCAGACCATCAACACCCAGCCCATCGACATCTCC	1363
1301	TTACTATTTTCATTGTCTCAGACAATCAATACTCAACCCATTGATATATCA	1350
1364	ACCGAGCTGAGCAAGGTGAACGCCTCCCTGCAGAACGCCGTGAAGTACAT	1413
1351	ACTGAGCTGAGTAAGGTTAATGCATCCCTCCAAAATGCCGTTAAATACAT	1400
1414	CAAGGAGAGCAACCACCAGCTGCAGAGCGTGAGCGTGAGCAGCAAGCGCC	1463
1401	AAAAGAGAGTAACCATCAACTCCAATCCGTTAGTGTAAGTTCTAAAAGAC	1450
1464	TGCACCGCGCCGCCATCTACACCGCCGAGATCCACAAGAGCCTGAGCACC	1513
1451	TTCATCGGGCAGCCATCTACACCGCAGAGATCCATAAAAGCCTCAGCACC	1500
1514	AACCTGGACGTGACCAACTCCATCGAGCACCAGGTGAAGGACGTGCTGAC	1563
1501	AATCTAGATGTAACATACTCAATCGAGCATCAGGTCAAGGACGTGCTGAC	1550
1564	CCCCCTGTTCAAGATCATCGGCGACGAGGTGGGCCTGCGCACCCCCCAGC	1613
1551	ACCACTCTTCAAAATCATCGGTGATGAAGTGGGCCTGAGGACACCTCAGA	1600
1614	GCTTCACCGACCTGGTGAAGTTCATCTCCGACAAGATCAAGTTCCTGAAC	1663
1601	GATTCACCTGACCTAGTGAAATTCATCTCTGACAAGATTAAATTCCTTAAT	1650
1664	CCCGACCGCGAGTACGACTTCCGCGACCTGACCTGGTGCATCAACCCCCC	1713
1651	CCGGATAGGGAGTACGACTTCAGAGATCTCACTTGGTGTATCAACCCGCC	1700
1714	CGAGCGGATCAAGCTGGACTACGACCAGTACTGCGCCGACGTGGCCGCCG	1763
1701	AGAGAGAATCAAATTGGATTATGATCAATACTGTGCAGATGTGGCTGCTG	1750
1764	AGGAGCTGATGAACGCCCTGGTGAACAGCACCCTGCTGGAGACCCGCACC	1813
1751	AAGAGCTCATGAATGCATTGGTGAACCTCAACTCTACTGGAGACCAGAACA	1800
1814	ACCAACCAGTTCCTGGCCGTGAGCAAGGGCAACTGCAGCGGCCCCACCAC	1863
1801	ACCAATCAGTTCCTAGCTGTCTCAAAGGGAAACTGCTCAGGGCCCACTAC	1850

1864	CATCCGGGGCCAGTTCAGCAACATGAGCCTGTCCCTGCTGGACCTGTACC	1913
1851	AATCAGAGGTCAATTCTCAAACATGTGCTGTCCCTGTTAGACTTGTATT	1900
1914	TGGGCCGGGGCTACAACGTGAGCAGCATCGTGACCATGACCAGCCAGGGC	1963
1901	TAGGTCGAGGTTACAATGTGTCATCTATAGTCACTATGACATCCCAGGGA	1950
1964	ATGTACGGCGGCACCTACCTGGTGGAGAAGCCCAACCTGAGCAGCAAGCG	2013
1951	ATGTATGGGGGAACCTACCTAGTGGAAAAGCCTAATCTGAGCAGCAAAAG	2000
2014	GAGCGAGCTGAGCCAGCTGAGCATGTACCGCGTGTTCGAGGTGGGCGTGA	2063
2001	GTCAGAGTTGTCACAACTGAGCATGTACCGAGTGTTTGAAGTAGGTGTTA	2050
2064	TCCGGAACCCCGGCCTGGGCGCCCCCGTGTTCACATGACCAACTACCTG	2113
2051	TCAGAAATCCGGGTTTGGGGGCTCCGGTGTTCATATGACAAACTATCTT	2100
2114	GAGCAGCCCGTGAGCAACGACCTGAGCAACTGCATGGTGGCCCTGGGCGA	2163
2101	GAGCAACCAGTCAGTAATGATCTCAGCAACTGTATGGTGGCTTTGGGGGA	2150
2164	GCTGAAGCTGGCCGCCCTGTGCCACGGCGAGGACAGCATCACCATCCCCT	2213
2151	GCTCAAACCTCGCAGCCCTTTGTACGGGGAAGATTCTATCACAATTCCT	2200
2214	ACCAGGGCAGCGGCAAGGGCGTGAGCTTCCAGCTGGTGAAGCTGGGCGTG	2263
2201	ATCAGGGATCAGGGAAAGGTGTCAGCTTCCAGCTCGTCAAGCTAGGTGTC	2250
2264	TGGAAGAGCCCCACCGACATGCAGAGCTGGGTGCCCCTGAGCACCGACGA	2313
2251	TGGAAATCCCCAACCGACATGCAATCCTGGGTCCCCTTATCAACGGATGA	2300
2314	CCCCGTGATCGACCGCCTGTACCTGAGCAGCCACCGCGGCGTGATCGCCG	2363
2301	TCCAGTGATAGACAGGCTTTACCTCTCATCTCACAGAGGTGTTATCGCTG	2350
2364	ACAACCAGGCCAAGTGGGCGCGTGGCCACCACCGCACCAGACAAGCTG	2413
2351	ACAACCAAGCAAAATGGGCTGTCCCGACAACACGAACAGATGACAAGTTG	2400
2414	CGCATGGAGACCTGCTTCCAGCAGGCCTGCAAGGGCAAGATCCAGGCCCT	2463
2401	CGAATGGAGACATGCTTCCAACAGGCGTGTAAGGGTAAATCCAAGCACT	2450
2464	GTGCGAGAACCCCGAaTGGGCCCCCCTGAAGGACAACCGCATCCCCAGCT	2513
2451	CTGCGAGAATCCCGAGTGGGCACCATTGAAGGATAACAGGATTCTTCAT	2500

[illegible]

65/73

Fig. 42B : F<sub>MU</sub>H<sub>MV</sub>-seq check: 4381 from: 1 to: 3126  
nucleic acid sequence of F<sub>MU</sub>H<sub>MV</sub>(non humanised)

ATGAAGGCTTTTCCAGTTATTTGCTTGGGCTTTGCAATCTTTTCATCCTC  
TATATGTGTGAATATCAATATCTTGCAGCAAATTGGATACATCAAGCAAC  
AGGTCAGGCAACTAAGCTATTACTCACAAAGTTCAAGCTCCTACGTAGTG  
GTCAAGCTTTTACCGAATATCCAACCCACTGATAACAGCTGTGAATTTAA  
GAGTGTAACCTCAATACAATAAGACCTTGAGTAATTTGCTTCTTCCAATTG  
CAGAAAACATAAACAATATTACGTCGCCCTCACCTGGGTCAAGACGTCAT  
AAACGGTTTGCTGGCATTGCCATTGGCATTGCgGCcCTCGGTGTTGCGAC  
CGCAGCACAAAGTGACTGCCGCTGTCTCATTAGTTCAAGCACAGACAAATG  
CACGTGCAATAGCAGCGATGAAAAATTCAATACAGGCAACTAATCGGGCA  
GTCTTCGAAGTGAAGGAAGGCACCCAACAGTTAGCTATAGCGGTACAAGC  
cATcCAAGACCATATCAATACTATTATGAACACCCAATTGAACAATATGT  
CTTGTCAGATCCTTGATAACCAGCTTGCAACCTCCCTAGGATTATACCTA  
ACAGAATTAACAACAGTGTTTCAGCCACAATTAATTAATCCAGCATTGTC  
ACCGATTAGTATACAAGCCTTGAGGTCTTTGCTTGGAAGTATGACACCTG  
CAGTGGTTCAAGCAACATTATCTACTTCAATTTCTGCTGCTGAAATACTA  
AGTGCCGGTCTAATGGAGGGTCAGATAGTTTCTGTTCTGCTAGATGAGAT  
GCAGATGATAGTTAAGATAAACGTTCCAACCATTTGTCACACAATCAAATG  
CATTGGTGATTGACTTCTACTCAATTTTCGAGTTTTATTAATAATCAAGAA  
TCCATAATTCAATTGCCAGACAGGATCTTGGAGATCGGAAATGAACAATG  
GCGCTATCCAGCTAAGAATTGTAAGTTGACAAGACACCACATATTCTGCC  
AATACAATGAGGCAGAGAGGCTGAGCCTAGAAACAAAACCTATGCCTTGCA  
GGCAATATTAGTGCCTGTGTGTTCTCATCTATAGCAGGGAGTTATATGAG

66/73

GCGATTTGTAGCACTGGATGGAACAATTGTTGCAAACGTGCGAAGTCTAA  
CGTGTCTATGCAAGAGTCCATCTTATCCTATATACCAACCTGACCATCAT  
GCAGTCACGACCATTGATCTAACGTCATGTCAAACATTGTCCCTGGACGG  
ACTGGATTTTCAGCATTGTCTCGCTAAGCAACATCACTTACGCTGAGAATC  
TTACTATTTTCATTGTCTCAGACAATCAATACTCAACCCATTGATATATCA  
ACTGAGCTGAGTAAGGTTAATGCATCCCTCCAAAATGCCGTTAAATACAT  
AAAAGAGAGTAACCATCAACTCCAATCCGTTAGTGTAAGTTCTAAAAGAC  
TTCATCGGGCAGCCATCTACACCGCAGAGATCCATAAAAGCCTCAGCACC  
AATCTAGATGTAACATACTCAATCGAGCATCAGGTCAAGGACGTGCTGAC  
ACCACTCTTCAAATCATCGGTGATGAAGTGGGCCTGAGGACACCTCAGA  
GATTCAGTACCTAGTGAAATTCATCTCTGACAAGATTAAATTCCTTAAT  
CCGGATAGGGAGTACGACTTCAGAGATCTCACTTGGTGTATCAACCCGCC  
AGAGAGAATCAAATTGGATTATGATCAATACTGTGCAGATGTGGCTGCTG  
AAGAGCTCATGAATGCATTGGTGAACATACTCTACTGGAGACCAGAACA  
ACCAATCAGTTCCTAGCTGTCTCAAAGGGAACTGCTCAGGGCCCACTAC  
AATCAGAGGTCAATTCTCAAACATGTCGCTGTCCCTGTTAGACTTGTATT  
TAGGTCGAGGTTACAATGTGTCTATAGTCACTATGACATCCCAGGGA  
ATGTATGGGGGAACTTACCTAGTGGAAAAGCCTAATCTGAGCAGCAAAAG  
GTCAGAGTTGTCACAACTGAGCATGTACCGAGTGTTTGAAGTAGGTGTTA  
TCAGAAATCCGGGTTTGGGGGCTCCGGTGTTCCATATGACAACTATCTT  
GAGCAACCAGTCAGTAATGATCTCAGCAACTGTATGGTGGCTTTGGGGGA  
GCTCAAACCTCGCAGCCCTTTGTACGGGGGAAGATTCTATCACAATTCCT  
ATCAGGGATCAGGGAAAGGTGTCAGCTTCAGCTCGTCAAGCTAGGTGTC  
TGGAATCCCCAACCGACATGCAATCCTGGGTCCCCTTATCAACGGATGA

67/73

TCCAGTGATAGACAGGCTTTACCTCTCATCTCACAGAGGTGTTATCGCTG  
ACAAcCAAGCAAAATGGGCTGTCCCGACAACACGAACAGATGACAAGTTG  
CGAATGGAGACATGCTTCCAACAGGCGTGTAAGGGTAAAATCCAAGCACT  
CTGCGAGAATCCCGAGTGGGCACCATTGAAGGATAACAGGATTCCTTCAT  
ACGGGGTCTTGTCTGTTGATCTGAGTCTGACAGTTGAGCTTAAAATCAAA  
ATTGCTTCGGGATTCGGGCCATTGATCACACACGGTTCAGGGATGGACCT  
ATACAAATCCAACCACAACAATGTGTATTGGCTGACTATCCCGCCAATGA  
AGAACCTAGCCTTAGGTGTAATCAACACATTGGAGTGGATACCGAGATTC  
AAGGTTAGTCCCTACCTCTTCAcTGTCCCAATTAAGGAAGCAGGCGAAGA  
CTGCCATGCCCCAACATACCTACCTGCGGAGGTGGATGGTGATGTCAAAC  
TCAGTTCCAATCTGGTGATTCTACCTGGTCAAGATCTCCAATATGTTTTG  
GCAACCTACGATACTTCCAGGGTTGAACATGCTGTGGTTTATTACGTTTA  
CAGCCCAgGCCGCTCATTTTtTTACTTTTATCCTTTTAGGTTGCCTATAA  
AGGGGGTCCCCATCGAATTACAAGTGGAATGCTTCACATGGGACCAAAAA  
CTCTGGTGCCGTCACCTTCTGTGTGCTTGCGGACTCAGAATCTGGTGGACA  
TATCACTCACTCTGGGATGGtGGGCATGGGAGTCAGCTGCACAGTCACCC  
GGGAAGATGGAACCAATCGCAGATAG

[SEQ ID NO: 97]

68/73

Fig 42C: F<sub>MUV</sub>humH<sub>M</sub>hum.seq check: 5778 from: 14 to: 3139  
Humanised nucleic acids sequence of F<sub>MUV</sub>H<sub>Mv</sub>

ATGAAGGCGTTCCCCGTGATCTGCCTGGGCTTCGCCATCTTCTCCAGCAG  
CATCTGCGTGAACATCAACATCCTGCAGCAGATCGGATACATCAAGCAGC  
AGGTGAGGCAGCTGAGCTACTACTCCCAGAGCTCCAGCTCCTACGTGGTG  
GTCAAGCTGCTGCCCAACATCCAGCCCACCGACAACAGCTGCGAGTTCAA  
GAGCGTGACCCAGTACAACAAGACCCTGAGCAACCTGCTGCTGCCCATCG  
CCGAGAACATCAACAACATCACCTCCCCCTCCCCCGGCTCCCGGGGGCAC  
AAGCGGTTCGCCGGCATCGCCATCGGCATCGCCGCCCTGGGCGTGGCCAC  
CGCCGCCCAGGTGACCGCCGCGGTGTCCCTGGTGCAAGGCCAGACCAACG  
CCCGCGCCATCGCCGCCATGAAGAACTCCATCCAGGCCACCAACCGCGCC  
GTGTTGAGGTGAAGGAGGGCACCCAGCAGCTGGCCATCGCCGTGCAGGC  
CATCCAGGACCACATCAACACCATCATGAACACCCAGCTGAACAACATGT  
CCTGCCAGATCCTGGACAACCAGCTGGCCACCTCCCTGGGCCTGTACCTG  
ACCGAGCTGACCACCGTGTTCCAGCCCCAGCTGATCAACCCCGCCCTGtc  
CCCCATCAGTATCCAGGCCCTGCGGTCCCTGCTGGGCAGCATGACCCCCG  
CCGTGGTGCAGGCCACCCTGAGCACCTCCATCAGCGCCGCCGAGATCCTG  
AGCGCCGGCCTGATGGAGGGCCAGATCGTGTCCGTGCTGCTGGACGAGAT  
GCAGATGATCGTGAAGATCAACGTGCCCACCATCGTGACCCAGTCCAACG  
CCCTGGTGATCGACTTCTACAGCATCAGCAGCTTCATCAACAACCAGGAG  
TCCATCATCCAGCTGCCCCGACCGCATCCTGGAGATCGGCAACGAGCAGTG  
GCGCTACCCCGCCAAGAACTGCAAGCTGACCCGCCACCACATCTTCTGCC  
AGTACAACGAGGCCGAGCGGCTGAGCCTGGAGACCAAGCTGTGCCTGGCC  
GGCAACATCAGCGCCTGCGTGTTCTCCAGCATCGCCGGCAGCTACATGCG

69/73

CCGCTTCGTGGCCCTGGACGGCACCATCGTGGCCAACTGCCGCAGCCTGA  
CCTGCCTGTGCAAGAGCCCCCTCCTACCCCATCTACCAGCCCGACCACCAC  
GCCGTGACCACCATCGACCTGACCTCCTGCCAGACCCTGAGCCTGGACGG  
CCTGGACTTCAGCATCGTGTCCCTGAGCAACATCACCTACGCCGAGAACC  
TGACCATCAGCCTGAGCCAGACCATCAACACCCAGCCCATCGACATCTCC  
ACCGAGCTGAGCAAGGTGAACGCCTCCCTGCAGAACGCCGTGAAGTACAT  
CAAGGAGAGCAACCACCAGCTGCAGAGCGTGAGCGTGAGCAGCAAGCGCC  
TGCACCGCGCCGCCATCTACACCGCCGAGATCCACAAGAGCCTGAGCACC  
AACCTGGACGTGACCAACTCCATCGAGCACCAGGTGAAGGACGTGCTGAC  
CCCCCTGTTCAAGATCATCGGCGACGAGGTGGGCCTGCGCACCCCCCAGC  
GCTTCACCGACCTGGTGAAGTTCATCTCCGACAAGATCAAGTTCCTGAAC  
CCCGACCGCGAGTACGACTTCCGCGACCTGACCTGGTGCATCAACCCCCC  
CGAGCGGATCAAGCTGGACTACGACCAGTACTGCGCCGACGTGGCCGCCG  
AGGAGCTGATGAACGCCCTGGTGAACAGCACCCCTGCTGGAGACCCGCACC  
ACCAACCAGTTCCTGGCCGTGAGCAAGGGCAACTGCAGCGGCCCCACCAC  
CATCCGGGGGCCAGTTCAGCAACATGAGCCTGTCCCTGCTGGACCTGTACC  
TGGGCCGGGGCTACAACGTGAGCAGCATCGTGACCATGACCAGCCAGGGC  
ATGTACGGCGGCACCTACCTGGTGGAGAAGCCCAACCTGAGCAGCAAGCG  
GAGCGAGCTGAGCCAGCTGAGCATGTACCGCGTGTTTCGAGGTGGGCGTGA  
TCCGGAACCCCGGCCTGGGCGCCCCCGTGTTCCACATGACCAACTACCTG  
GAGCAGCCCGTGAGCAACGACCTGAGCAACTGCATGGTGGCCCTGGGCGA  
GCTGAAGCTGGCCGCCCTGTGCCACGGCGAGGACAGCATCACCATCCCCT  
ACCAGGGCAGCGGCAAGGGCGTGAGCTTCCAGCTGGTGAAGCTGGGCGTG  
TGGAAGAGCCCCACCGACATGCAGAGCTGGGTGCCCTGAGCACCGACGA


70/73

CCCCGTGATCGACCGCCTGTACCTGAGCAGCCACCGCGGCGTGATCGCCG  
ACAACCAGGCCAAGTGGGCCGTGCCCACCACCCGCACCGACGACAAGCTG  
CGCATGGAGACCTGCTTCCAGCAGGCCTGCAAGGGCAAGATCCAGGCCCT  
GTGCGAGAACCCCGAaTGGGCCCCCCTGAAGGACAACCGCATCCCCAGCT  
ACGGCGTGCTGAGCGTGGACCTGAGCCTGACCGTGGAGCTGAAGATCAAG  
ATCGCGAGCGGCTTCGGCCCCCTGATCACCCACGGCAGCGGCATGGACCT  
GTACAAGAGCAACCACAACAACGTGTACTGGCTGACCATCCCCCCCATGA  
AGAACCTGGCCCTGGGCGTGATCAACACCCTGGAGTGGATtCCCCGCTTC  
AAGGTGAGCCCCTACCTGTTACCGTGCCCATCAAGGAGGCCGGCGAGGA  
CTGCCACGCCCCGACCTACCTGCCCCGCCGAGGTGGACGGCGACGTGAAGC  
TGAGCAGCAACCTGGTGATCCTGCCCCGGCCAGGACCTGCAGTACGTGCTG  
GCCACCTACGACACCAGCCGCGTGGAGCACGCCGTGGTGTACTACGTGTA  
CAGCCCCGGCCGCAGCTTCTTCTACTTCTACCCCTTCCGCCTGCCCATCA  
AGGGCGTGCCCATCGAGCTGCAGGTGGAGTGCTTCACCTGGGACCAGAAG  
CTGTGGTGCCGCCACTTCTGCGTGCTGGCCGACAGCGAGAGCGGCGGCCA  
CATCACCCACAGCGGCATGGTGGGCATGGGCGTGAGCTGCACCGTGACCC  
GCGAGGACGGCACCAACCGCCGCTAG

[SEQ ID NO: 98]

**Fig. 43** Total IgG directed against the FHN protein


71/73



SUBSTITUTE SHEET (RULE 26)

**Fig. 44** IgG1, directed against the FHN protein.








72/73



SUBSTITUTE SHEET (RULE 26)

**Fig. 45** IgG2a directed against the FHN protein

73/73



SUBSTITUTE SHEET (RULE 26)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int .tional Application No

PCT/EP 00/01903

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19709512 A	10-09-1998	NONE	
WO 9103552 A	21-03-1991	US 5166057 A AT 126272 T AU 636916 B AU 6411890 A CA 2065245 A DE 69021575 D DE 69021575 T DK 490972 T EP 0490972 A ES 2075901 T GR 90100639 A JP 5500607 T PT 95124 A US 5252289 A US 6001634 A US 5578473 A US 5854037 A US 5840520 A US 5786199 A US 5820871 A ZA 9006852 A	24-11-1992 15-08-1995 13-05-1993 08-04-1991 01-03-1991 14-09-1995 14-12-1995 30-10-1995 24-06-1992 16-10-1995 30-12-1991 12-02-1993 18-04-1991 12-10-1993 14-12-1999 26-11-1996 29-12-1998 24-11-1998 28-07-1998 13-10-1998 31-07-1991
WO 9610641 A	11-04-1996	EP 0704533 A AU 3607695 A EP 0783586 A FI 971272 A NZ 293600 A	03-04-1996 26-04-1996 16-07-1997 26-05-1997 28-01-1999

# INTERNATIONAL SEARCH REPORT

Int .tional Application No

PCT/EP 00/01903

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PALESE P. ET AL.: "Negative-strand RNA viruses: Genetic engineering and applications." PROC. NATL. ACAD. SCI. U.S.A., vol. 93, October 1996 (1996-10), pages 11354-11358, XP000196755 page 11354, right-hand column, last paragraph -page 11356, right-hand column, paragraph F	5
P,X	NEUMANN G: ET AL.: "Plasmid-driven formation of influenza virus-like particles." JOURNAL OF VIROLOGY, vol. 74, no. 1, January 2000 (2000-01), pages 547-551, XP002140119 ISSN: 0022-538X the whole document	1,3-5, 12, 18-23, 25-31,33
P,A	FLICK R. AND HOBOM G.: "Interaction of influenza virus polymerase with viral RNA in the 'corkscrew' conformation." JOURNAL OF GENERAL VIROLOGY, vol. 80, no. 10, October 1999 (1999-10), pages 2565-2572, XP002140120 ISSN: 0022-1317 figure 1	7-11

# INTERNATIONAL SEARCH REPORT

Int. J. Application No

PCT/EP 00/01903

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ZHOU Y. ET AL.: "Membrane-anchored incorporation of a foreign protein in recombinant Influenza virions." VIROLOGY, vol. 246, 20 June 1998 (1998-06-20), pages 83-94, XP002110226 the whole document	5
A	ZOBEL A. ET AL.: "RNA polymerase I catalysed transcription of insert viral cDNA." NUCLEIC ACIDS RESEARCH, vol. 21, no. 16, 1993, pages 3607-3614, XP002110227 page 3607, right-hand column, paragraph 2 page 3612, right-hand column, paragraph 2 -page 3613, left-hand column, line 1 page 3614, left-hand column, paragraph 2	15-17
A	WO 96 10641 A (BAYER AG ; HOBOM GERD (DE); NEUMANN GABRIELE (DE); MENKE ANNETTE (D) 11 April 1996 (1996-04-11) cited in the application the whole document	6-11
A	FLICK R. ET AL.: "Promoter elements in the influenza vRNA terminal structure." RNA, vol. 2, no. 10, 1996, pages 1046-1057, XP000914725 ISSN: 1355-8382 the whole document	6-11
A	NEUMANN G. AND HOBOM G.: "Mutational analysis of influenza virus promoter elements in vivo." JOURNAL OF GENERAL VIROLOGY 1995, vol. 76, no. 7, 1995, pages 1709-1717, XP002140118 ISSN: 0022-1317 cited in the application	6-11
A	PICCONE M. E. ET AL.: "MUTATIONAL ANALYSIS OF THE INFLUENZA VIRUS vRNA PROMOTER" VIRUS RESEARCH, vol. 28, no. 2, 1 January 1993 (1993-01-01), pages 99-112, XP000619019 ISSN: 0168-1702 the whole document	6-11

-/--

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01903

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/86 C12N7/01 C12N5/10 A61K39/00 A61K39/145  
A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE 197 09 512 A (HOBOM GERD PROF DR DR) 10 September 1998 (1998-09-10)  the whole document	1, 4, 5, 12, 14, 18-33
Y	WO 91 03552 A (SINAI SCHOOL MEDICINE) 21 March 1991 (1991-03-21)  figure 11; example 7	1, 4, 5, 12, 14, 18-33
Y	TAKASE H. ET AL: "Antibody responses and protection in mice immunized orally against influenza virus." VACCINE, vol. 14, no. 17/18, 1996, pages 1651-1656, XP002110225 page 1652, left-hand column, paragraph 1	27

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 June 2000

Date of mailing of the international search report

07/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Mandl, B



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/45, 15/62, C07K 14/115, 14/12, 14/135, A61K 39/155, 39/165</b>		<b>A3</b>	(11) International Publication Number: <b>WO 00/18929</b>
			(43) International Publication Date: 6 April 2000 (06.04.00)
(21) International Application Number: <b>PCT/EP99/07004</b>			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 20 September 1999 (20.09.99)			
(30) Priority Data: 9820931.5 25 September 1998 (25.09.98) GB 9906868.6 24 March 1999 (24.03.99) GB			
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM BIOLOGICALS S.A. [BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BOLLEN, Alex [BE/BE]; Université Libre de Bruxelles, Faculté des Sciences, Rue de l'Industrie 24, B-1440 Nivelles (BE). HOUARD, Sophie [BE/BE]; CRI - Centre de Recherches Industriel/ULB, Département de Biologie Moléculaire, Rue de l'Industrie 24, B-1400 Nivelles (BE).			
(74) Agent: PRIVETT, Kathryn, Louise; Corporate Intellectual Property, SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			Published With international search report.
			(88) Date of publication of the international search report: 9 November 2000 (09.11.00)
(54) Title: <b>PARAMYXOVIRUS VACCINES</b>			
(57) Abstract  Heterochimeric proteins or immunogenic derivatives thereof are described comprising immunogenic fragments of RSV, PIV1, PIV2, PIV3, MV and MuV fusion and attachment glycoproteins. Such heterochimeric proteins may be expressed, in particular, in CHO cells and may be used in vaccine compositions to treat respiratory disorders such as those caused by paramyxoviridae viral antigens.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Licchtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## INTERNATIONAL SEARCH REPORT

International Application No

PC., EP 99/07004

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 7 C12N15/45 C12N15/62 C07K14/115 C07K14/12 C07K14/135  
 A61K39/155 A61K39/165

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal, BIOSIS, MEDLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HOMA FL ET AL.: "Development of a novel subunit vaccine that protects cotton rats against both human respiratory syncytial virus and human parainfluenza virus type 3" JOURNAL OF GENERAL VIROLOGY, vol. 74, no. 9, September 1993 (1993-09), pages 1995-1999, XP002134292 READING GB cited in the application figure 1	3-8, 16-20, 22-29
X	WO 93 14207 A (CONNAUGHT LAB) 22 July 1993 (1993-07-22) cited in the application examples 9-15	3-8, 16-20, 22-29



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the international search

28 July 2000

Date of mailing of the international search report

17.08.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Cupido, M

## INTERNATIONAL SEARCH REPORT

International Application No

PC, EP 99/07004

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	OIEN NL ET AL: "Vaccination with a heterologous respiratory syncytial virus chimeric FG glycoprotein demonstrates significant subgroup cross-reactivity" VACCINE, vol. 11, no. 10, October 1993 (1993-10), pages 1040-1048, XP002143652 GUILDFORD GB the whole document	3-20, 22-29
A	WO 94 25600 A (SMITHKLINE BEECHAM BIOLOG; BOLLEN ALEX (BE); HOUARD SOPHIE (BE); N) 10 November 1994 (1994-11-10) cited in the application the whole document	1-29
A	LEHMAN JD ET AL.: "Comparison of soluble and secreted forms of human parainfluenza virus type 3 glycoproteins expressed from mammalian and insect cells as subunit vaccines" JOURNAL OF GENERAL VIROLOGY, vol. 74, no. 3, March 1993 (1993-03), pages 459-469, XP002134293 READING GB cited in the application page 463, last paragraph -page 465	1-29
A	LELIÈVRE D ET AL.: "Structural properties of chimeric peptides containing a T-cell epitope linked to a fusion peptide and their importance for in vivo induction of cytotoxic T-cell responses" EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 249, no. 3, 1 November 1997 (1997-11-01), pages 895-904, XP000929575 the whole document	1-29

# INTERNATIONAL SEARCH REPORT

In' ational application No.  
PCT/EP 99/07004

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claim 28 is directed to a method of treatment of the human or animal body, the search has been carried out and based on the alleged effects of the vaccine.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

As a result of the prior review under R. 40.2(e) PCT,  
no additional fees are to be refunded.

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☒ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/EP 99/07004

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-29 (all partially)

Heterochimaeric protein comprising an immunogenic fragment of the fusion (F) protein of RSV, process for preparing it, DNA encoding said protein, vectors and host cells containing this DNA and corresponding vaccines and uses thereof.

2. Claims: 1-12, 16-29 (all partially)

Heterochimaeric protein comprising an immunogenic fragment of the fusion (F) protein of PIV1, process for preparing it, DNA encoding said protein, vectors and host cells containing this DNA and corresponding vaccines and uses thereof.

3. Claims: 1-12, 16-29 (all partially)

Heterochimaeric protein comprising an immunogenic fragment of the fusion (F) protein of PIV2, process for preparing it, DNA encoding said protein, vectors and host cells containing this DNA and corresponding vaccines and uses thereof.

4. Claims: 1-29 (all partially)

Heterochimaeric protein comprising an immunogenic fragment of the fusion (F) protein of PIV3, process for preparing it, DNA encoding said protein, vectors and host cells containing this DNA and corresponding vaccines and uses thereof.

5. Claims: 1-12, 16-29 (all partially)

Heterochimaeric protein comprising an immunogenic fragment of the fusion (F) protein of MV, process for preparing it, DNA encoding said protein, vectors and host cells containing this DNA and corresponding vaccines and uses thereof.

6. Claims: 1-29 (all partially)

Heterochimaeric protein comprising an immunogenic fragment of the fusion (F) protein of MuV, process for preparing it, DNA encoding said protein, vectors and host cells containing this DNA and corresponding vaccines and uses thereof.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT, EP 99/07004

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9314207 A	22-07-1993	AU 3340293 A	03-08-1993
		CA 2126863 A	22-07-1993
		EP 0621898 A	02-11-1994
		FI 943211 A	02-09-1994
		JP 7501707 T	23-02-1995
		NO 942530 A	05-09-1994
		RU 2123047 C	10-12-1998
		US 6033668 A	07-03-2000
		US 5968776 A	19-10-1999
		US 5998169 A	07-12-1999
		US 6017539 A	25-01-2000
WO 9425600 A	10-11-1994	AU 6964894 A	21-11-1994
		CA 2161645 A	10-11-1994
		CN 1126492 A	10-07-1996
		EP 0695358 A	07-02-1996
		JP 8512199 T	24-12-1996